

John Fotheringham, Dr. C. R. Sneath, Dr. J. N. Anderson, Dr. R. J. Dwyer, Dr. Gibb Wishart, Dr. S. Johnston, Dr. D. Anderson, Dr. H. Glendenning, Dr. P. G. Goldsmith, Dr. Charles Trow, Dr. N. A. Powell, Dr. J. L. Bradley, Dr. W. B. Bryan, Dr. Andrew Eadie, Dr. E. H. Greene, Dr. C. H. Thomas, Dr. H. Clare, Dr. J. McConnell, Dr. J. Rowan, Dr. George Strathy, Dr. J. A. Kinnear, Dr. H. Becker, Dr. M. J. Perkins, Dr. A. E. McKay, Dr. A. G. A. Fletcher, Dr. C. B. Shuttleworth, Dr. W. A. Peart, Dr. E. Clouse, Dr. W. B. Kendal, Dr. C. A. Simple, Dr. J. M. McCormack, Dr. M. O. Scott, Dr. A. H. Garratt, Dr. C. A. Campbell, Dr. M. A. Kendrick, Dr. Brefney, O'Reilly, Dr. M. L. G. McGibbon, Dr. Walter McKeown, Dr. W. H. Lowry, Dr. W. S. Harrison, Dr. A. Jukes Johnson, Dr. R. B. Nevitt, Dr. N. Anderson, Dr. J. K. M. Gordon, Dr. H. C. Parsons, Dr. C. P. Lusk, Dr. Reade, Dr. T. H. Ashley, Dr. Henwood, Dr. A. Davidson, Dr. F. C. Stephenson, Dr. John E. King, Dr. G. W. Brand, Dr. Thomas Mason, Dr. Burgess, Dr. F. Fenton, Dr. J. T. Clarke, and Messrs. Albert Crux, M. L. Kendrick, J. L. Bingham, and S. T. White.—Report in *Mail and Empire*.

---

Do not operate for foreign body in the knee joint without first excluding dislocation of one of the semilunar cartilages.

FRACTURES of the neck of the femur in old people sometimes cause no other symptoms than disability. The mildness of the trauma and the freedom from much pain should not deceive one.

IN amputations below the knee, insist on active and passive motion in the knee joint at an early date. If this is not done contracture ensues, which makes the application of an artificial limb difficult.

IN compound fractures involving loss of continuity do not needlessly remove any piece of bone that has even the smallest attachment. It is surprising how often such pieces heal into the wound and thereby help to save loss of substance.

IN cases of renal colic do not make too positive a pre-operative diagnosis of calculus, no matter how typical the symptoms may be. It has happened very often at the time of operation that no stone is found. Fortunately, these cases are nearly always cured by the exploratory nephrotomy.—*American Journal of Surgery*.