

Deer Park, Lower Arrow Lake, B.C.; Revelstoke, B.C.;
Sicamous, B.C. (*John Macoun.*)

GALIUM APARINE, L.; Macoun, Cat. Can. Plants, Vol. I,
p. 200.

The only western station for this plant given by Prof. Macoun is Victoria, Vancouver Island. It has since been found to be of very general distribution in British Columbia. Our specimens are from Elk River Bridge, Rocky Mountains. (*Dr. Geo. M. Dawson.*) Deer Park, Lower Arrow Lake, Columbia River, B.C.; Ainsworth, Kootanie Lake, B.C.; Kamloops, B.C.; Yale, B.C.; Agassiz, B.C.; Lulu Island, B.C.; Salt Spring Island, Gulf of Georgia; common on Vancouver Island. (*John Macoun.*) Specimens from some of the above localities have been distributed as var. *Vaillantii*, Koch, but they are true *G. Aparine*. We agree with Dr. Greene that though this species is "as much at home in our woods and thickets as any indigenous plant, it is probable that it came hither from the Old World within the last two centuries."

SHERARDIA ARVENSIS, L.

In fields, Victoria, Vancouver Island, 1893. (*John Macoun.*) Not before recorded west of Ontario.

ASPERULA ARVENSIS, L.

Edge of a marsh, near Hamilton, Ont. (*J. M. Dickson.*)
New to Canada.

LIATRIS SCARIOSA, Willd.; Macoun, Cat. Can. Plants,
Vol. I., p. 208.

Near Sarnia, Lambton Co., Ont. (*C. K. Dodge.*) Not recorded from Ontario since collected by MacLagan. Not having seen MacLagan's specimens, Prof. Macoun, when preparing his Catalogue, was of the opinion that MacLagan's stations for *L. scariosa* should be referred to *L. cylindrica*.