

*Anaphora mortipennella*, Grote ♂. Labial palpi reflexed, thrown back over and as long as the dorsum of thorax, but not closely applied, thickly scaled but less so than in allied species, fuscous outwardly along the sides, dead whitish on the inside. Head and thorax above dead or dirty whitish. Primaries pale, dirty whitish, with heavily sprinkled black scales on costal region at base, fading outwardly. A black scale patch at extremity of discal cell, and a larger one on submedian fold, below median vein, at about the middle of the wing; parallel with this at base, a few black scales. There is a faint sprinkling of black scales over the median nervules, and about internal angle are two or three better marked black points on the margin. Four costal black marks before the apex, the first of these above discal spot; other costal marks towards the base of the wing. Fringes fuscous, faintly lined. Secondaries fuscous, much darker than, and strangely contrasting with, the pallid primaries. Beneath both wings fuscous with ochrey stains. The basal joint of labial palpi is prominently dark fuscous or blackish outwardly. *Expanse* 25 *m. m.* Central Alabama. June.

Smaller than *A. flumifrontella*, and easily recognized by its pallid discoloured fore wings, which are also a little more determinate at apices and internal angle than usual.

*Anaphora agrotipennella*, Grote ♂. Fuscous or blackish wood brown. Labial palpi reflexed, and as long as the dorsum of thorax, a little paler inwardly, blackish outwardly. Primaries above fuscous, blackish, with a light purplish reflection. From the base outwardly, below median vein, is a prominent pale streak fading externally, where it is diffuse and dark ochrey. It is bordered beneath at base by black scales like a dash, and surmounted and partly interrupted by a black scale patch below median nervure before vein 2. On the discal cell is an unprominent black scale patch towards the base, beyond which an obscure ochrey longitudinal median shade, sometimes lost, stretches over the nervules, and is interrupted at the extremity of the cell by a distinct black subquadrate scale patch. Faint blackish costal and terminal marks; fringes fuscous. Secondaries and their fringes fuscous. Beneath, both wings and body parts blackish-fuscous. *Expanse* 27 *m. m.* Central Alabama; June and July. Very common.

I have only seen males of this species, in which the ornamentation of the fore wings above recalls that of various species of *Agrotis*, such as *A. jaculifera*, etc. I have tried to recognize in this species *A. Poppeanella*,