

Under side : palpi, head, thorax, legs and abdomen light brown. On the fore wings the ground colour is light brown ; the fascial bar that is mentioned above is here very prominent ; the apical area has a large dark brown triangular spot. On the hind wings the ground colour is the same as on the fore wings ; the markings are the same as in *T. eurycles*, Lat., of Brazil. Fringes on both pairs of wings are a shade lighter than the ground colour. Tails,  $\frac{3}{4}$  inch in length. Expanse,  $1\frac{7}{8}$  inch. Hab.: San Pedro Sula, Honduras, Central America.

This species seems to be very rare, for of the many hundreds of specimens that I have obtained from Dr. C. Thieme, of Honduras, it remains unique in my collection.

*Thymele Borja*, n. sp.—Upper side : antennæ, head, thorax and abdomen are very dark brown. On both pairs of wings the ground colour is of a beautiful fawn-brown ; margins edged with dark brown ; fringes a shade lighter than the ground colour.

Under side : palpi light brown ; antennæ, thorax, legs and abdomen dark brown. Fore wings lighter than above ; transparent spots the same as above ; there are two dark wavy brown lines on the fascia. On the hind wings the ground colour has the same shade of brown as the fore wings ; on the costa near the base there is a quadrate spot ; aside from this there are two dark brown bars on the fascia. Tails,  $\frac{7}{16}$  inch long. Expanse, 2 inches. Hab.: Barja, Bolivia, S. America.

*Goniurus Triptolemus*, n. sp.—Female. Upper side : antennæ, head, thorax and abdomen dark brown. On the fore wings the ground colour is also dark brown ; on the fascia there is a narrow transparent bar, which is composed of four unequal spots ; the third spot from the costa is the largest. In the median cell beyond this bar is a narrow transparent spot which crosses the cell ; on the subapical space is a row of four very minute transparent spots ; fringes brown. On the hind wings the ground colour is dark brown, but a shade lighter than the fore wings. The tails are white ; the vein that extends through them is sprinkled with brown. Apical fringes are white, but slightly intermixed with brown. Fringes on the outer margin and on the tails are very long and pure white. Abdominal fringes are shorter and brown.

Under side : the thorax and abdomen are much lighter than above ; the abdomen has a dark brown bar ; underneath the legs are also dark brown. On the fore wings the ground colour is lighter than above ; all the same markings, but the transparent bar is more sharply defined ; the inner margin is pale brown, and dusted with white. On the hind wings the ground colour is the same as the fore wings ; there is a faint dark