Figure No.		Page No.
34.	Geographic distribution of pH levels measured in Adirondack lakes higher than 610 meters elevation, June 24-27, 1975.	3–90
35.	Susceptibility map of Quebec lake waters.	3-93
36.	Sensitivity to acidification of surface waters in the Maritime provinces.	3–95
37.	Sensitivity to acidification of surface waters in insular Newfoundland.	3–96
38.	Varying effects of lake pH on the distribution of mercury in ecosystems.	5–3
39.	Seasonal and spatial distribution of long-term trends in average airport visibilities for the eastern United States.	5-10
40.	Effects of fine particle increments on calculated visual range.	5- ₁₂
41.	Light-scattering per unit mass of sulphate aerosol as a function of relative humidity.	5-13
A1	Bedrock classification for eastern United States showing sensitivity to acidification.	A-5
A2	Acid precipitation sensitivity - Bedrock and soils derived from bedrock.	A-11
A3	Sensitivity of glacial and postglacial surficial sediments to acid loading.	A-13
A4	Map of counties with soils judged to be sensitive to acid precipitation in the eastern United States.	A-15
A5	Regions with significant areas of soils that are non sensitive, slightly sensitive and sensitive with the eastern U.S.	A-17
A6 ·	Relative sensitivity of soils of eastern Canada to acid rain	. A-20
A7	Acid precipitation overlay of cropland in the eastern United States.	A-22
A8	Relative terrain sensitivity to acid precipitation	A-25
A9	Ecodistrict sensitivity to acidification for forest productivity and aquatic inputs for the province of Ontario.	A-29
A10	Sensitivity of soils to acid precipitation.	A-31
A11	Sensitivity of vegetation to SO ₂ .	A-32
A12	Combined terrain sensitivity to acid precipitation.	A-33

J