

again resected ; this time five inches, going well within the border of the attached mesentery, united the ends with the Murphy button, ligatured the mesenteric vessels and brought the mesenteric borders together with catgut sutures close up to the bowel. There were thus 11 inches of bowel removed. The hernial sac was excised, and the peritoneal wound closed with mattress sutures of silk. The muscular borders were next closed with buried sutures of silk-worm gut and the pectineal fascia was attached to the re-united Poupart's ligament by three sutures of catgut. Finally the skin was closed by a separate layer of silk-worm gut sutures, and a small tent of iodoform gauze introduced at the lower angle of the wound. The operation lasted two hours and was well borne. The patient never had a bad symptom and made an uneventful recovery. A liquid motion, (with flatus), was expelled at the end of twenty-four hours, (after administration of an enema). A regular movement occurred again next day, and on the fifth, sixth, ninth and tenth days. The button was found imbedded in a well formed stool, which was passed at 1.30 p.m., October 30th, just ten full days after operation. The wound was perfectly healed and the patient allowed up on the 22nd. Healing per primam.

My experience in these three cases leads me to the conclusion that the Murphy button is a valuable aid in end to end anastomosis of intestine. So many artificial aids have been introduced for this purpose, have had their day and have been discarded, that most surgeons are now sceptical about anything of this kind. It is, of course, not to be assumed that union of intestine cannot be secured without such aids, for it undoubtedly can, but the great desiderata, rapidity of operation and accuracy and security of co-aptation, are both admirably effected by this instrument. I cannot agree with the view which has recently been promulgated, that the Murphy button is useful in the hands of the tyro and is not necessary to the experienced surgeon. The actual union of the intestinal ends is but one part of the operation, even if it be the culminating point,