

- Face tuberculate or arched, not carinate. 47
47. Thorax and scutellum very densely pollinose above.
- PTERALLASTES
- Scutellum not densely pollinose above. 48
48. Third joint of antennæ broad, thorax not vittate. 49
- Third joint oval; thorax usually vittate above. HELOPHILUS
49. Thickly pilose. MALLOTA
- Thinly pilose. 50
50. Hind tibiæ of male with an internal median spur.
- TEUCHOCNEMIS
- Hind tibiæ of male not with an internal median spur. POLYDONTA
51. Hind femora thickened, with a bifid spur below on outer end.
- STENOGASTER
- Hind femora not with a bifid spur below. 52
52. Slender species, abdomen narrowed on basal portion. BACCHA
- Abdomen in no wise club-shaped or narrowed basally. 53
53. Thorax with distinct yellow markings other than on the humeri. 62
- Thorax not with distinct yellow markings other than rarely on the humeri. 54
54. Face transversely arched or carinate, not produced, not tuberculate; abdomen more or less elongate and nearly bare. 55
- More or less thickly pilose species, often large. 57
55. Hind femora extraordinarily thickened; anterior cross-vein rectangular. SYRITTA
- Hind femora more or less thickened; cross-vein oblique. 56
56. Face sharply carinate. TROPIDIA
- Face transversely arched. XYLOTA
57. Scutellum, margin of the thorax, and the pleuræ with bristly hairs. CHRYSOCHLAMYS
- Thorax wholly without bristles. 58
58. Face short, not produced, concave from antennæ to tip, not tuberculate; hind femora thickened. 59
- Face produced, long. 60
59. Abdomen elongate. BRACHYPALPUS
- Abdomen very broad, thorax densely pilose, very large species.
- POCOTA