An irregular winter visitor, never abundant. Breeds in the far north.

240. Cedar Waxwing.

(Ampelis cedrorum.)

A resident species, common, and generally distributed in summer, irregular in winter, its movements at that season probably depending upon the food supply.

FAMILY LANIIDÆ. SHRIKES.

GENUS LANIUS.

241. Northern Shrike.

(Lanius borealis.)

A regular, but not abundant winter visitor. Breeding far north. Arrives in October, departs in April.

242. Loggerhead Shrike.

(Lanius Iudovicianus.)

A regular, but not abundant summer resident, breeds throughout its range in the Province.

Our form is probably referable to subspecies L. l. migrans. Arrives early in April, departs about the end of August.

FAMILY VIREONIDÆ. VIREOS.

GENUS VIREO.

SUBGENUS VIREOSYLVA.

243. Red eyed Vireo.

(Vireo olivaceus.)

A common summer resident in woods. Breeds throughout its range in the Province.

Arrives early in May, departs towards the end of September.

244. Philadelphia Vireo.

(Vireo philadelphicus.)

Tolerably common summer resident, breeds throughout its range in the Province.

Arrives about the middle of May, departs at the end of September.

245. Warbling Vireo.

(Vireo gilvus,)

Common summer resident, breeds throughout its range in the Province.