

- KILLDEER, *Oxyechus vociferus*—Common.
- RUFFED GROUSE, *Bonasa umbellus*—Found in swampy places.
- MARSH HAWK, *Circus hudsonius*—Not very common.
- SHARP-SHINNED HAWK, *Accipiter velox*—One seen some distance up country.
- RED-TAILED HAWK, *Buteo borealis*—A pair bred in a big woods half a mile from camp.
- SPARROW HAWK, *Falco sparverius*—A few seen up country.
- OSPREY, *Pandion haliaetus carolinensis*—One fishing near Bobcaygeon, August 12th, 1912.
- KINGFISHER, *Ceryle alcyon*—Very common.
- HAIRY WOODPECKER, *Dryobates villosus*—Common.
- DOWNY WOODPECKER, *Dryobates pubescens*—Common.
- YELLOW-BELLIED SAPSUCKER, *Syphrapicus varius*—Rather numerous. Breeds.
- RED-HEADED WOODPECKER, *Melanerpes erythrocephalus*—A number at the Point. Much more numerous up country.
- FLICKER, *Colaptes auratus*—A few observed each summer. Not common.
- NIGHTHAWK, *Chordeiles virginianus*—Flocks of fifteen or twenty seen during migrations.
- CHIMNEY SWIFT, *Chatura pelagica*—A few always flying around.
- HUMMINGBIRD, *Archilochus colubris*—One or two seen each year.
- KINGBIRD, *Tyrannus tyrannus*—Quite common and as usual very noisy.
- CRESTED FLYCATCHER, *Myiarchus crinitus*—To be seen or heard every day.
- PHOEBE, *Sayornis phæbe*—Plasters nest under the rocky ledges in Fenelon river.
- OLIVE-SIDED FLYCATCHER, *Nuttallornis borealis*—This rare species is one of the features of Pleasant Point bird life. It is quite a common breeder and may be seen almost any time of the day sitting on the top of some dead tree watching for passing insects.
- WOOD PEWEE, *Myiochanes virens*—Heard calling from the deep woods.
- ALDER FLYCATCHER, *Empidonax trailli alnorum*—Heard one calling from a swampy thicket.
- LEAST FLYCATCHER—*Empidonax minimus*—Fairly common.