

7. *costotristrigella*. *N. sp.*

Head and palpi pale saffron yellow, the outer surface of the palpi dark brown; antennae dark brown; thorax and basal portion of the dorsal margin of the primaries dark brown dusted with yellowish white, the primaries otherwise white dusted with dark brown, with an oblique dark costal streak near the base, extended to the fold; just before the middle is another longer one also extended to the fold where it enlarges into an irregular spot, being also dusted with yellowish white above the fold; just behind the middle is another streak not reaching the fold, behind which is a small costal brown spot and a row of brown spots around the apex. The apical portion of the wing is more densely dusted than the disc. Ciliae white with fuscous spots. *Alar ex.* $\frac{1}{2}$ inch. Kentucky, in August and September. Taken flying.

8. *T. bimaculella*. *N. sp.*

Outer surface of the second joint of the labial palpi brown; inner surface and terminal joint pale yellowish or stramineous; head pale stramineous; antennae pale yellowish tinged with fuscous; thorax shining dark brown, almost black, except the tip which is stramineous; costal half of the primaries fuscous, narrow towards the base, but spreading towards the apex, where it is mixed with pale yellowish, with a distinct dark brown spot beyond the end of the disc; dorsal half stramineous, widest at the base, narrowing towards the apex, with a distinct dark brown spot within the margin about the middle: ciliae fuscous and stramineous mixed; anterior and intermediate legs dark brown, the tarsi faintly annulate with stramineous; posterior legs stramineous. *Alar ex.* $\frac{1}{3}$ inch. Kentucky.

9. *T. aurosuffusella*. *N. sp.*

Palpi pale stramineous, the outer surface of the second joint of the labial pair brown; antennae pale fuscous; head and thorax pale stramineous, with a small pale fuscous spot on the thorax before the apex; primaries pale stramineous streaked along the fold with pale reddish golden, and the apical portion of the wing suffused with the same hue; a rather wide pale fuscous streak, the basal portion of which is scalloped towards the fold, extends from the base of the costa along the costal margin to a little beyond the middle, and a similar streak, scalloped towards the fold in its posterior half, extends along the dorsal margin from near the base to the