

The specimen in Mr. Belanger's collection differs from all of my specimens in having the dorsal spot prolonged towards the apex till it meets in the apical part of the wing with the second costal streak, in having the apical spot more definite, and the dusting in the ciliæ so arranged as to form two short hinder marginal lines. I have no doubt it is the same species. The pattern of ornamentation (not the colors) of this species is almost exactly that of the *right* wing in Mr. Stainton's figure of *B. demaryella*, but the left wing in the figure is very different. *Vid. Nat. Hist. Tin., v. 7.*

*Lithocolletis salicifoliella*, CAN. ENT., v. 3., p. 163.

“ *Scudderella* Frey. & Boll. ? *Ent. Zeitung*, p. 212.

In the collection received from Mr. Belanger is a single specimen of this species, together with the mined leaf of poplar, from which it was bred. It has heretofore been found only in Willow leaves. It is an exceedingly variable species, some specimens being so densely dusted with brown or gray brown scales as to suggest a resemblance to the European species *L. populifoliella*, whilst others are almost free from dusting, and may be described as having a ground color of pale golden or saffron marked with white. In the former description of this species I was led by the resemblance of the more densely dusted specimens to *L. populifoliella*, as figured by Mr. Stainton in *Nat. Hist. Tin., v. 2*, and the position in which *L. pastorella* is there placed in his classification of the species of this genus, to suggest that it might be identical with *pastorella*. Perhaps the following description may convey a better idea of an average specimen than the one previously given

Thorax and primaries bright golden or saffron yellow, according to the light, or even sometimes dull brownish yellow, the thorax and basal portion of the dorsal margin of the fore wings being largely intermixed with white and dusted more or less with black; sometimes the inner angle is of the general ground hue, scarcely dusted or marked with either white or dark brown, and then there is a median white basal streak which meets at an acute angle with a dorsal white streak about the basal fourth of the wing length. Both of these white streaks and all other white markings on the wings are more or less dusted with dark gray brown, sometimes so much as to obscure the white. Before the middle of the costa is a long white streak, which attains the middle of the wing, curving backwards; a little behind this, on the dorsal margin, is a large dorsal white streak, wide