

shoulder at the sound side, and carry it round the back of the neck, so as to be visible at the opposite side, where it is to be held fast, place the point behind the elbow of the injured arm and draw down the end in front of the patient. Next bend the arm carefully and place it across the chest on the middle of the cloth. Then take the lower end upwards over the shoulder on the wounded side, and knot to the other end at the nape of the neck. This done draw the point forward round the elbow, and fasten it with a pin. This is called the larger arm sling.

**WOUND ON THE CHEST.**—Place the middle of the bandage on the chest with the point over the shoulder, carrying the two ends round the chest and knot at the back. Next draw the point over the shoulder downwards and tie or pin it to one of the ends.

**WOUND OF THE HAND.**—Take a bandage, spread it out, and lay the wrist on the lower border with the fingers towards the point. Next turn the point over the fingers and carry it over the wrist. This done take the ends round the wrist, fixing the point, cross them, carry them back again and knot together. Take a second bandage and support the forearm in the larger sling as above.

**WOUND OF THE HIP.**—Fold one bandage narrow, and tie it round the body as a waist-belt above the hips. Lay the centre of a bandage on the wound with the point upwards, pass the ends round the upper part of the thigh, cross and carry it to the front and knot them together, next pass the point under the waist-belt and fasten it with a pin.

**WOUND OF THE FOOT.**—Take a bandage, spread it out and place the sole of the foot on its centre, with the toes in the direction of the point. Draw the point upwards over the toes and instep of the foot; then take the ends forward above the ankles, and cross in front of the leg, carry them downwards under the sole of the foot, and knot them together above the ankle.

**TO SECURE FRACTURES.**—Surgical or improvised splints may be adjusted to a broken limb by taking two triangular bandages folded broad or narrow, according to circumstances, and tying them securely one above and the other below the fracture. As many more bandages can be added as may be considered necessary to *secure* the limb.

**FRACTURED COLLAR BONE.**—Place a pad in arm-pit on injured side, and suspend the arm in a large arm sling, then fix the arm to the side with a small narrow bandage passing round the chest and fastened under the sound arm.

The triangular bandage may be applied in many other ways: but the above directions are quite sufficient to indicate its different uses.