

1582. *L. officinale*, L. (Common Gromwell.)

Introduced and very abundant in rocky pastures. Ju—2. (B).

1585. *L. hirtum*, Lehm. (Hairy Puccoon.)

River side in sand. On the banks of the Ottawa above Aylmer.

Rare. (Mrs. Chamberlin) June—3. Flowers bright orange, woolly inside.

SYMPHYTUM, L. Comfrey.

1590. *S. officinale*, L. (Common Comfrey.)

Introduced. Rideau rifle range. Buckingham. Aylmer. July—2.

LYCOPSIS, L. Bugloss.

1591. *L. Arvensis*, L. (Small Bugloss.)

Introduced. Rare in sandy fields. Theodore street and at Billings Bridge. June—2.

ECHIUM, L. Viper's Bugloss.

1592. *E. vulgare*, L. (Blue weed.)

Introduced. Gradually becoming a troublesome weed. Ju—2.

CONVOLVULACEÆ. Bindweed Family.

IPOMÆA, L. Morning Glory.

1594. *I. purpurea*, L. (Common Morning Glory.)

A garden escape, frequently found in waste places.

CONVOLVULUS, L. Bindweed.

1597. *C. SPITHAMÆUS*, L. (Low Bindweed.)

Calystegia spithamea, Pursh.

Clay banks, sandy shores, and rocky islands. Rare. Bank of the Ottawa below Rockcliffe. Kettle Island. and formerly on the rocky island opposite the end of Bank street. A very attractive plant with short stems and large pure white flowers. July—1.

1598. *C. sepium*, L. (Bracted Convolvulus. Hedge Bindweed.)

Calystegia sepium, R. Br.

Introduced. Chelsea Road, one locality. July—1. This is evidently the European form, which has in some way got introduced. It climbs up over high bushes fifteen or twenty feet from the ground and has snow-white flowers.