

Territories with their chief characters.—Continued.

Time of seeding.	Colour, Size, Arrangement of Flowers and other Characters of the Plant.	Method of Propagation and Distribution.	Place of Growth, and Products Injured.	Methods of Eradication.
g. Sept.	Whitish, 1-in., spikes peduncled; pods oblong, 1-in., covered with hooked prickles.	Seeds, pods attached to stock, in hay.	Summer fallows, pastures; wool.	Summer fallow early.
y. Aug.	Yellow, 1-in., leafy cymes; leaves 3-parted; whole plant dark green, hairy.	" "	Summer fallows, grain fields.	Summer fallow, cultivate.
y. Sept.	Bright yellow, 1-in., solitary on long stalks.	Seeds, runners	Lowlands, particularly if alkaline.	" "
r. Sept.	Pink to deep rose, 2½ in. corymb....	" rootstocks	Summer fallows, grain fields.	Summer fallow early, harrow, cultivate.
Sept.	Yellow, 1½ in., open at night, leafy spike.	Seeds, wind..	Summer fallows.	Pull, plough fall and spring.
Sept.	White, turning pink, 2-in., malodorous leafy spikes; buds nodding; lower glistering white, simple, branching at the top.	Seeds and rootstocks.	Grain fields on knolls.	Summer fallow, cultivate thoroughly.
.....	White; umbel 1-in. across; stem stout, spotted with purple, strong smelling, very poisonous.	Seeds, carried by floods.	Wet meadows, troublesome in hay, poisonous to stock.	Spud, mow in flower.
.....	Red, much bearded inside, 1-in. dense terminal and axillary spikes; berry reddish, 1-in.	Seeds and running rootstocks.	Nowly broken land summer fallows and pastures.	Break early, summer fallow.
.....	White, small, in large terminal panicles.	" "	Grain fields, pastures.	Summer fallow, cultivate.
Sept.	Bright yellow; 1-in.; whole plant glutinous; but bearing large drops of white resin.	Seeds, wind, in hay.	Fields, pastures, road sides.	Mow, cultivate.
.....	Yellow, head large, 1-sided.....	Seeds, running rootstocks, wind.	Grain fields, summer fallows.	Plough and cultivate.
Oct.	White; 1-in., crowded on spreading branches.	" "	Grain fields, sandy land.	" "
Sept.	White; heads very numerous, small, crowded in a slender erect wand-like panicle.	Seeds, wind...	Summer fallows, grain fields.	Summer fallow early, cultivate fall and spring.
Oct.	Inconspicuous; 1 in., hanging, short-stalked in axils of the upper leaves; leaves less than 1-in., rough, oblong linear, entire, opposite below, alternate above.	Copious underground creeping stems.	Grain fields.....	Summer fallow, cultivate constantly.
ov.	Green, 1 in., crowded in large terminal panicles; stem smooth.	Seeds, wind, floods.	Grain crops, road sides.	Pull, mow.
"	Yellow, 1 in., sterile flowers in terminal racemes or spikes, fertile flowers axillary at base of spikes; stems rough; seed 1 in., bearing a crown of 8-10 tubercles above the middle; leaves 3-lobed.	Seeds, in grain, wind, floods.	Low rich land, grain fields, wheat.	Pull, mow, burn old plants.
"	Yellow, 1 in., sterile in racemes, fertile green, axillary; seed 1 in., long, hairy without spines. Plant grayish-green.	Seeds, running rootstocks.	Rich cultivated land, all crops.	Summer fallow early, cultivate late.