

*Tyrrell.) Miles Canon, Lat. 62°, north of British Columbia, 1887. (Dr. G. M. Dawson. Wm. Ogilvie.)*

*PHACELIA SERICEA*, Gray; Macoun, Cat. Can. Plants, Vol. I., pp. 333 and 567.

*P. sericea*, Gray, var. *Lyallii*, Gray; Macoun, Cat. Can. Plants, Vol. I., p. 333 in part and p. 567.

Common in the Rocky Mountains from Lat. 52° south to the International Boundary, Roger's Pass, Selkirk Mountains, alt. 4,500 ft.: Mount Arrowsmith, Vancouver Island, alt. 5,500 ft. (*John Macoun.*)

*PHACELIA SERICEA*, Gray, var. *LYALLII*, Gray.

Our only Canadian specimens of this species were collected in 1895 by Prof. Macoun on the high slopes of Sheep Mountain, Waterton Lake, Rocky Mountains.

*AMSIANKIA ECHINATA*, Lehm.

Near an old mining camp at Revelstoke, B.C., July 21st, 1890. Introduced from the United States. (*John Macoun.*) New to Canada.

*AMSIANKIA LYCOPSOIDES*, Lehm.

*A. lycopsoides*, Lehm. var. *bracteosa*, Macoun, Cat. Can. Plants, Vol. I., p. 568.

Mary Island, Gulf of Georgia, B.C., 1885. (*Dr. G. M. Dawson.*) Our only Canadian specimens. - The type was collected by Scouler on an island or along the shores of the Straits of Juan de Fuca.

*AMSIANKIA LYCOPSOIDES*, Lehm., var. *BRACTEOSA*, Gr.; Macoun, Cat. Can. Plants, Vol. I., p. 338.

Common on Vancouver Island, B.C.

*MYOSOTIS COLLINA*, Hoffm.

Low fields, Edmonton, Ont., 1890. (*Jas. White.*) New to Canada. Introduced.