

Watson, Proc. Amer. Acad. 1873, viii. 600.— *O. densiflora* var. *tenella* Gray, Proc. Amer. Acad. 1873, viii. 384.— *Boisduvalia Torreyi* Watson, Bot. Calif. 1876, i. 233; Index, 363.— A. foot to a foot and a half high, slender, simple or with a few long erect virgate branches chiefly toward the base, densely villous, often hoary; leaves commonly under 1 in. long, narrowly lanceolate, acute, entire or minutely denticulate, the upper shorter and not widened; flowers axillary along the branches, about equaling the leaves; corolla 2 to 4 mm. long, violet; capsules slender, generally curved outwards, tapering above, more or less 2-edged, about 10 mm. long, loculicidal, the lateral nerves commonly heavier; seeds 6 to 8 in each cell, similar to those of *densiflora* but smaller, about .5×1 mm. (varying from .38 to .63×.88 to 1.6 mm.).— Washington to Idaho and Central California.

Specimens examined from Washington (Brandegge, 1882 and 1883, 788; Seattle, Piper, 1888, 411; Spokane, Henderson, 1892, 2468 in part, and Sandberg, Aug. 1892; Klickitat Co., Suksdorf, Aug. 8, 1881; Falcon Valley, Suksdorf, Aug. 1880, 378, and July 3, 1882), Oregon (Henderson, 1882, 360; Howell, Aug. 1880; Hall, 1871, 189; Grant's Pass, Howell, June 23, 1884; Idaho (Kootenai Co., Heller, 1892, 905), and California (Rockville, Earle, Aug. 1, 1880; Yreka, Greene, 1876, 853; Humboldt & Sonoma Counties, Bolander, 1866-7, 6535; Almaden, Torrey, 1865, 109; Borax Lake and Bear Mountain, Torrey, 1865; Sta. Lucia Mts., Vasey, July 1880, Kernville, Coville and Funston, 1891, 1041; Lake Co., Brandegge, July 1884, and June 1890; Howell Mt., Brandegge, Aug. 1888; Plumas Co., Brandegge, July 4, 1892; Byron Sta., Greene, May 24, 1886; Newcastle, Brandegge, July 5, 1889; Sonoma, Brandegge, June 1892; Modoc Co., Mrs. Austin, Aug. 1885; Humboldt Co., Mrs. Bush.)

B. DENSIFLORA (Lindley) Watson, Bot. Calif. 1876, i. 223; Index, 362.— *B. Douglasii* Spach, Hist. Nat. Végétaux, 1835, iv. 385, atlas, pl. 85, f. 2; Monographia