

- Obstetrical Society of London, 195
 Occiput, 255, 265
 backward rotation of, 267
 forward rotation of, 267
 Occurrence, in shoulder presentation,
 341
 of eclampsia, complicating labour,
 443
 Oidium albicans, 567
 Oldham's perforator, 673
 Oligo-hydramnios, 139
 Olshausen, 473
 Oosperm, 9
 Ophthalmia neonatorum, 581
 causes of, 581
 symptoms and treatment, 582
 Oponins, 529
 Organs, genital, involution of, during
 puerperium, 489
 Os uteri, see *Uterus, os of*
 Outlet of pelvis, 223, 231
 Over-distension of lower uterine seg-
 ment, 406, 414, 515
 Ovulation, 1
 and menstruation, 6
 during pregnancy, 74
 Ovum, abnormal conditions of, 95
 abnormal implantation of, 95
 after escape from Graafian
 follicle, 3
 apoplectic, 128
 complete, 27
 complete, from about the ninth
 week, 44
 complete, from the fourth or fifth
 week, 43
 complete, of three months' de-
 velopment retained for some
 weeks after death, 210
 earliest described, 12
 "external wandering" of, 4
 fertilisation and embedding of, 9
 forceps, 209
 Graf Spee's sagittal section of, 24
 in earliest stages of development,
 12, 15
 Leopold's, 25, 35
 mamalian, 19
 of four weeks' gestation, 20
 Peters', 12, 15, 18, 35
 segmentation of, 11
 Teacher and Bryce's, 15, 16, 24
- PAJOT's manoeuvre, 632
 Palfyn's forceps, 624
 Palpation, abdominal, in diagnosis of
 face presentations, 311
 in diagnosis of normal labour,
 281, 283-286
 Paralysis, birth, 579
 facial, 579, 580
 Parametritis, remote, 534
- Paré, Ambroise, 612
 Parietal obliquity, 258, 261
 Parturition, see *Labour*
 Passages, maternal, in first and second
 stages of labour, 225
 abnormal conditions of bony
 pelvis, 308
 abnormal conditions of soft parts,
 308
 Pasteur and puerperal fever, 504
 Pasteurisation of milk, 559
 Pedicle, abdominal or ventral stalk,
 24
 Pelvimeter, Collin, 363
 Skutsch, 368
 Pelvimetry, clinical, 363
 external, 364
 internal, 366
 Pelvis æquabiliter justo-major, 358
 minor, 358
 Pelvis, axis of, 234
 bisected, 228
 brim of, outline in principal
 varieties of contracted pelvis,
 391
 elliptical or simple flat, 358
 false, 226
 female, 225, 227
 flat, 358, 360, 361, 371, 373
 floor of, 232, 399
 generally enlarged, 362
 kyphotic, 388, 390
 ligature or excision of veins of,
 for pyæmia, 540
 local inflammation in, 532
 Malacostean, 383
 male, 226
 measurements of, 363-366
 Nägele's, 386
 non-rachitic, 358
 normal, 230, 365
 oblique, due to disease of hip-joint,
 388
 of a woman who died five days
 after delivery, frozen section,
 492
 of a woman who died immediately
 after delivery, frozen section,
 487
 osteomalacic, 383
 pseudo-osteomalacic, 385
 rachitic flat, 358-361
 reniform flat, 358
 Robert's, 387
 spondylolisthetic, 389, 390
 triradiate, 383
 true, 226-237
 veins in puerperal infection, 515
 Pelvis, contracted, 357
 due to disease affecting skeleton
 generally, 383
 due to disease of pelvic joints, 386
 due to disease of hip-joints, 389