

Walkerville, Ontario, providing, of course, the edition has not previously been exhausted.

ENDOMETRITIS.

By J. J. O'SULLIVAN, M.D., New York City.

Being a friend of Glyco-Thymoline for many years, I have no hesitancy in endorsing it at any time. As regards my experience with it in Gynecology, will say that I have a record of some ninety cases in which I have used Glyco-Thymoline to a greater or less extent, and have always found it of great value in reducing engorgements and promoting a healthy condition of the tissues. The following cases serve to illustrate the usual method followed in applying this agent:

Case 1.—Mrs. H. G., aged 24, married three years, multipara, occupation, housewife; gave the following history: Begun menstruating at the age of 14 years and had always been very regular during the menstrual period, had always had some pain which had, however, become intense during the last two. Past two months had suffered severe back-ache and pain throughout the pelvic region; bowels constipated. Digital examination showed the cervix to be very tender and engorged with some slight congestion of the uterus itself, accompanied by a profuse whitish discharge. Diagnosis of cervicitis being made, a tampon of cotton soaked in pure Glyco-Thymoline was applied and patient directed to inject small amount of Glyco-Thymoline pure into vagina twice a day. Tampons of cotton and Glyco-Thymoline were repeated every other day and patient discharged in one month cured. Aside from an occasional laxative, no other treatment was used.

Case 2.—Mrs. M. H., aged 21, married; multipara; occupation, housewife; came to me complaining of intense pain throughout the pelvic region, feeling of weight and bearing down sensation; bowels constipated, and frequent micturition, having to void her urine from five to six times nightly which was accompanied by severe burning and tenesmus. An examination disclosed a lacerated cervix with considerable inflammation of the endometrium. Treatment consisted of tampons of cotton and pure Glyco-Thymoline applied every second day with intra-uterine douches of a hot 25 per cent. solution of Glyco-Thymoline applied by means of Chamberlain's glass tube. This patient has been under my treatment for three months now, and the laceration has almost healed, which I expect to be complete in two or three weeks when I will discharge her, the inflammation of the endometrium having long since disappeared.