

SUMMARY OF KEY GREENHOUSE GASES AFFECTED BY HUMAN ACTIVITIES					
	Carbon Dioxide	Methane	CFC-11	CFC-12	Nitrous Oxide
Atmospheric concentration	ppmv	ppmv	pptv	pptv	ppbv
Pre-industrial (1750-1800)	280	0.8	0	0	288
Present day (1990)	353	1.72	280	484	310
Current rate of change per year	1.8 (0.5%)	0.015 (0.9%)	9.5 (4%)	17 (4%)	0.8 (0.25%)
Atmospheric lifetime (years)	(50-200)†	10	65	130	150

ppmv = parts per million by volume;

ppbv = parts per billion (thousand million) by volume;

pptv = parts per trillion (million million) by volume.

† The way in which CO₂ is absorbed by the oceans and biosphere is not simple and a single value cannot be given; refer to the main report for further discussion.

Two important greenhouse gases, water vapour and ozone, are not included in the table above. Water vapour has the largest greenhouse effect, but its concentration in the troposphere is determined internally within the climate system, and, on a global scale, is not affected by human sources and sinks. Water vapour will increase in response to global warming and further enhance it; this process is included in climate models. The concentration of ozone is changing both in the stratosphere and the troposphere due to human activities, but it is difficult to quantify the changes from present observations.

For a thousand years prior to the industrial revolution, abundances of the greenhouse gases were relatively constant. However, as the world's population increased, as the world became more industrialized and as agriculture developed, the abundances of the greenhouse gases increased markedly. The figures below illustrate this for carbon dioxide, methane, nitrous oxide and CFC-11.

Since the industrial revolution the combustion of fossil fuels and deforestation have led to an

increase of 26% in carbon dioxide concentration in the atmosphere. We know the magnitude of the present day fossil-fuel source, but the input from deforestation cannot be estimated accurately. In addition, although about half of the emitted carbon dioxide stays in the atmosphere, we do not know well how much of the remainder is absorbed by the oceans and how much by terrestrial biota. Emissions of chlorofluorocarbons, used as aerosol propellants, solvents, refrigerants and foam blowing agents, are also well known; they were not present in the atmosphere before their invention in the 1930s.

The sources of methane and nitrous oxide are less well known. Methane concentrations have more than doubled because of rice production, cattle rearing, biomass burning, coal mining and ventilation of natural gas; also, fossil fuel combustion may have also contributed through chemical reactions in the atmosphere which reduce the rate of removal of methane. Nitrous oxide has increased by about 8% since pre-industrial times, presumably due to human activities; we are unable to specify the sources, but it is likely that agriculture plays a part.