

Phthisis, General medicinal treatment of (continued)

- Iodoform, II. 8-9, 12
- Menthol, II. 14
- Myrtol, II. 14
- Turpentine, II. 13
- Arsenic in, II. 24-25
- Cacodylate of sodium in, II. 25
- Calmette's ophthalmic-reaction in, II. 21
- Classification of phthisis according to amenability to treatment, II. 1-3
- Cod-liver oil in, II. 27-28
- Conditions favourable to cure of, II. 3-4
- Iron in, II. 26-27
- Maragliano's antitoxic serum in, II. 22
- Mineral waters in, II. 28-31
- Serum treatment of, II. 22, 23
- Tuberculin in, II. 16-21
- See also* Additional Formulæ, II. 31-32

Phthisis, Indications and prophylaxis in, I. 715-25

- Avoidance of unhealthy occupation, I. 720
- Catarrhal attacks to be guarded against, I. 720
- Dangers of puberty and of the period of rapid development, I. 719
- Emaciation in, I. 719-20
- Indications for treatment of, I. 715
- Influence of heredity, I. 716-17
- Management of infants and children with phthisical tendencies, I. 717-18
- Measures for preventing the dissemination of infective germs, I. 722-25
- Pulmonary gymnastics, I. 720
- Relationship between acid and phthisis, I. 721
- Risk of re-infection, I. 722
- Tendency to scrofulous affections, I. 719

Phthisis, Laryngeal, II. 62-66

- Bismuth subnitrate in, II. 64
- Chinosol in, II. 64
- Chloroform in, II. 66
- Cocaine in, II. 64, 66
- Diet in, II. 65
- Dysphagia in, II. 64
- Eucaïne in, II. 64, 66
- Formalin in, II. 65
- Inhalations in, II. 63, 65, 66
- Insufflations in, II. 64, 66
- Intratracheal injections in, II. 65
- Iodine in, II. 63
- Iodoform in, II. 64
- Lactic acid in, II. 65
- Menthol in, II. 64

Phthisis, Laryngeal (continued)

- Morphine in, II. 64, 66
- Orthoform in, II. 64, 66
- Sprays in, II. 63

Phthisis, Regiminal Treatment of, I. 726-38

- Air and ventilation, I. 731-32
- Alcohol in, I. 729
- "Alimentation forcée," I. 729
- Bodily exercise, I. 733
- Clothing, I. 735
- Cold douches and sponging, I. 734-35
- Cream in, I. 728
- Daily dietary in, I. 730-31
- Farinaceous foods in, I. 729
- Hydrotherapy in, I. 734
- Importance of regiminal and hygienic treatment of, I. 726, 735
- Koumiss in, I. 728
- Pneumatic treatment of, I. 732-33
- Rest in, I. 732
- Sanatorium treatment in, I. 726, 736-38
- Value of fats and carbohydrates, I. 727

Phthisis, Symptomatic treatment of, II. 33-60

- OF COUGH, II. 41-47**
 - Alkaline drinks in, II. 43
 - Belladonna in, II. 45
 - Bromide of potassium in, II. 44
 - Butyl chloral in, II. 44
 - Chloral hydrate in, II. 44
 - Cigarettes in, II. 47
 - Codeina in, II. 45
 - Counter-irritation in, II. 33-34, 44
 - Heroin in, II. 45, 47
 - Inhalations in, II. 43, 47
 - Ipecacuanha lozenges in, II. 46
 - Morphine in, II. 46
 - Nitrate of silver in, II. 46
 - Opium in, II. 42, 45
 - Sodium bromide in, II. 44
 - Syrups in, II. 46
 - Terpine hydrate in, II. 47
- OF DIARRHŒA, II. 58-60**
 - Bismuth carbonate in, II. 59
 - Bismuthi subgallas in, II. 59
 - Castor oil in, II. 59
 - Coto bark in, II. 60
 - Creasote in, II. 59
 - Enema in, II. 59
 - Kino in, II. 59
 - Lime water in, II. 59
 - Liquor calcis saccharatus in, II. 59
 - Nitrate of silver in, II. 60
 - Opium in, II. 59
 - Salicylate of bismuth in, II. 59
 - Sulphate of copper in, II. 60