

What Clinical Trial Shows

- 1. Duration of the disease is cut short.
- The most marked and disagreeable symptoms are ameliorated.
- 3. Temperature is reduced.
- 4. Delirium is terminated or much abated.
- 5. Tympanites is almost wholly avoided.
- 6. Diarrhea is checked.

remarkable an powerful, in the capable of destroying any discesse-germ with which it can be brought into contact; safe, in that it may be taken internally, in aqueous solution, in quantities limited



only by the capacity of the patient to swallow and absorb it.

Supplied in ounce, half-ounce and quarter-ounce bottles, also in vials of 15 grains, six vials in a box.

Full Literature Sent Free on Request

For Diagnosis of Typhoid Fever

Our Typhoid Agglutometer simplifies the Widal test, obviating the use of the microscope and the fresh live culture of typhoid bacilli. It is


fully equal to the former method in delicacy and accuracy, and much superior in convenience.

The apparatus consists of four tubes of a sterile permanent suspension of dead typhoid bacilli, a tube

of blood-serum diluent, an empty tube for collecting the blood, a needle for puncture,

and a pipette for withdrawing the separated serum. It contains material for filteen to thirty tests.

Directions Accompany Each Package Circular Free



PARKE. DAVIS & COMPANY

LABORATORIES: DETROIT, MICH., U.S.A.; WALKERVILLE, ONT.; HOUNSLOW, ENG.

Branches: New York, Chicago, St. Louis, Boston, Baltimore, New Orleans, Kansas City, Indianapolis,
Minneapolis, Memphis; London, Eng.; Montreal, Que.; Syoney, N.S.W.;
St. Petersburg, Russia: Bimla, India; Tokio, Japan.