

A. Stewart, Palmerston, Ont.; W. T. Connell, Kingston, Ont.; T. H. Stark, Toronto; A. Stark, Berwick, Ont.; J. M. Jory, St. Catharines, Ont.; J. W. S. McCullough, Alliston, Ont.; E. Meek, Port Rowan, Ont.; J. H. Mullin, Hamilton, Ont.; J. F. Uren, Toronto; A. H. Halliday, Traverse City, Mich.; O. J. McCully, Monkton, Ont.; George Elliott, Toronto; R. J. Trimble, Queenston, Ont.; W. S. Harrison, Toronto; W. H. Harris, Toronto; W. O. Stewart, Guelph, Ont.; J. T. Hart, Toronto; F. Winnett, Toronto; E. Sisley, Maple, Ont.; A. R. Gordon, Toronto; E. H. Stafford, Toronto; A. T. Macnamara, Toronto Junction; D. H. Hogg, London, Ont.; A. T. Hobbs, London, Ont.; E. T. Snider, Brussels, Ont.; C. E. Stacey, Toronto; L. Bentley, Toronto; C. H. Thomas, Gormley, Ont.; Beattie Nesbitt, Toronto; Chas. Trow, Toronto; R. H. Green, Embro, Ont.; R. N. Fraser, Thamesville, Ont.; Thos. Kerr, Toronto; Wm. Kerr, Cayuga, Ont.; B. Spencer, Toronto; D. W. Gordon, Lucknow, Ont.; J. P. Kennedy, Wingham, Ont.; W. J. O. Malloch, Toronto; W. H. Groves, Burnhamthorpe, Ont.; H. H. Sinclair, Walkerton, Ont.; D. B. Bentley, Sarnia, Ont.; A. W. McFaul, Stayner, Ont.; D. King Smith, Toronto; J. H. Cotton, Toronto; F. Montizambert, Ottawa; Hugh Bain, Prince Albert, N.W.T.; J. H. Elliott, Gravenhurst, Ont.; J. Montgomery, Oshawa, Ont.; Dr. Hodgkin, Deer Park, Ont.; Julia Thomas, Toronto; Dr. Jackes, Eglington, Ont.; W. H. Clemes, Toronto; W. T. Sherris, Toronto; R. O. Snider, Toronto; Ernest Hall, Toronto; W. J. McCollum, Toronto; J. H. Henwood, Toronto; A. G. Ashton Fletcher, Toronto; B. Z. Milner, Toronto; W. J. Wilson, Toronto; Graham Chambers, Toronto; Lelia H. Davis, Toronto; C. J. O. Hastings, Toronto; Geo. H. Carveth, Toronto; Fred. Feinton, Toronto; H. A. Bruce, Toronto; C. L. Starr, Toronto; and J. P. Russell, Toronto.

TUBERCULOSIS IN CANADIAN CATTLE AND ITS PREVENTION.

Dr. J. GEORGE ADAMI (Montreal).—At the outset, he stated that there were three questions to be asked and answered: 1. Is tuberculosis in cattle a source of danger to other cattle, so as to seriously affect their well-being, and to be a source of loss to the owners? 2. If infectious from animal to animal, is it infectious from animal to man, and thereby a grave source of danger to the human race? 3. If infectious from animal to man, what are the commonest modes of infection, and, as a sequel to this, how are we to diminish the danger? If the first can be answered in the affirmative, how can the disease be erased? To do this we should employ all the means in our power. What organization and official steps should be taken in our country to stamp it out? In regard to the question: "Is it dangerous to other cattle when in cattle?" there is abundant evidence to show that the introduction of an infected bull into a