

BRACHIOPODA.

RHYNCHONELLA PSITTACEA. (Gm.) Owen.

Henley Harbor, 3 to 8 fathoms; Temple Bay, 10 fathoms;
Dead Island, near Square Island, 1 to 4 fathoms.

*List of Labrador shells recorded by Dr. A. S. Packard, Jr.,
but not obtained by the Stearns Expedition.*

GASTROPODA.

Bela cancellata (Migh.) Stimp.


Fusus syrtensis, nov. sp. (Not seen.)

Fusus tornatus Gld. (Neptunca despecta (L.) Ad.).

Buccinum tenue Gray.

? Natica heros Soub. (Lamatia heros (Say) H. & A. Ad.).

Littorina palliata Gld.




Scalaria Grœnlandica Perry.

Margarita campanulata Morse, nov. sp.

(M. helicina, var.).

Scissurella crispata Flem.

Adeorbis costulata Stimp. (Molleria
costulata (Moll.) Jeff., Fr.).? Lamellaria perspicua Loren. (Not
seen.)Bulla pertenuis Migh. (Diaphana per-
tenuis (Migh.) Verrill).Bulla occulta Migh. (Cylichna occulta
(Migh.) H. & A. Ad.).

Dendronotus arborescens Ald. & Han.

PTEROPODA.

Fig. 1. *Kennerlia glacialis*. x 2. Specimen from
Labrador, collected by Prof.
A. S. Packard, Jr., 1864.

Cylichna limacina Phipps (*C. papiliona-*
eca Pallas).

Limacina helicina Phipps.

Fig. 1 a. The same. Inter-
ior view of another specimen.

LAMELLIBRANCHIATA.

Solen ensis Linne (*Ensatella Americana* (Gould) Verrill).