

CHENOPODIUM BOSCIANUM, Moq.

Sandy thickets, Pelee Point, Lake Erie. No. 54,724.
(John Macoun.) New to Canada.

AQUILEGIA COLUMBIANA, Rydb., Bull. Torr. Bot. Club, XXIX: 145.

This species is somewhat intermediate between *A. formosa* and *A. truncata* and has been mistaken for both. It has the habit, spur and sepals of the former and the short truncate lamina of the latter. From Banff to Alaska.

DELPHINIUM BROWNII, Rydb., Bull. Torr. Bot. Club, XXIX: 148.

This is most nearly related to *D. glaucum* but differs in the puberulent leaves with narrow segments, the lax raceme with more erect pedicels and the darker flowers. Described from specimens collected at Banff in 1893 by Addison Brown. A common plant in that region.

RANUNCULUS ALLENI, Robinson, Rhodora, VII: 220.

R. affinis var. *leiocarpus*, Cat. Can. Pl. 1: 18.

First collected by Mr. J. A. Allen on Mt. Albert, Gaspé, Que., in 1881, the next year by John Macoun at the same place, No. 1,015, and in 1883 on Table Top Mountain not far from Mt. Albert by James Porter, No. 68,678. Other localities cited by Dr. Robinson are: Okkak, Labrador, and Rama, Labrador.

RANUNCULUS GLABERRIMUS, Hook.

Pentt. ton, Lake Okanagan, B.C., April 12th. 1903. No. 59,519. (*W. Spreadborough*.) Not recorded from that region.

RANUNCULUS YUKONENSIS, Britt.

Near Peace River Landing, Atha., No. 59,521. June 8th, 1903. (*J. M. Macoun*.) Recorded before only from the Yukon district.

CAULOPHYLLUM THALICTROIDES, Mx.

In woods along the Assiniboine River near Portage La Prairie, Man. In flower, May 31st, 1906. (*W. Herriot*.) Western limit in Canada