

The Canadian Entomologist.

VOL. XXV.

LONDON, OCTOBER, 1893.

No. 10.

WASHINGTON TENTHREDINIDÆ AND UROCERIDÆ.

BY ALEX. D. MACGILLIVRAY, ITHACA, N. Y.

The specimens on which the present paper has been based were collected by Mr. Trevor Kincaid, of Olympia, Washington. They were all collected near that city except a few, which were taken along the Skokomish River. Great credit is due Mr. Kincaid for bringing together so large a collection. The specimens are deposited in the entomological collection of Cornell University.

TENTHREDINIDÆ.

Trichiosoma † *triangulum*, Kirby.

3 ♂ ♂, June 6, 22, 23, 1892. They agree perfectly with Norton's description of this species, except the apical segments of the antennæ, which are entirely black.

Zarea americana, Cress.

2 ♀ ♀, April 17 and May 23, 1892.

Hylotoma abdominalis, Leach.

1 ♀, May 3, 1892. Skokomish River.

Hylotoma Macleayi, Leach.

1 ♀, May 11, 1892.

Euura abbiricta, Cress.

1 ♀, Skokomish River, May 8, 1892.

Nematus castaneus, Kirby.

A single female, which in all probability belongs to this species, has two broad black bands on the lateral lobes of the mesothorax and two less distinct bands on the median lobe; the square spot on the vertex is fuscous, a small spot between the antennæ and the base of the sinus from

† The following is omitted from Cresson's synopsis. It will probably prove to be a variety of *triangulum* and not the European species.

vitellina, Linn.

1882. *Trichiosoma*, Kirby, List Hymen., Brit. Mus., I., 70.

Habitat.—Vancouver's Island, Rocky Mountains, (Europe).