

*Lycæna rusticus*.—As many as four eggs in succession on the upper side of the erect leaves of *Dodecatheon meadia* var. *lancifolium*.

*Anthocharis ausonides*.—Mostly underneath the leaves of *Arabis* [? sp.].

*Colias Behrii*.—In the heart of the flower buds or tender leaves of *Vaccinium cæspitosum* and *Gentiana Newberryi*.

*Parnassius smintheus*.—On the side or top or hidden out of sight on the leaves of *Phlox cæspitosa*, *Carex filifolia*, *Gayophytum diffusum*, or on pine burrs, sticks and rocks.

*Papilio eurymedon*.—Under the leaves of *Ceanothus prostratus*.

*Pamphila sabuleti*.—At or near the base of the stem of *Carex filifolia* and under the leaves of *Trifolium monandrum*.

*Pyrgus cæspitalis*.—In the centre of the plant, *Horkelia fusca*.

*Eudamus tityrus*.—On the underside of the leaves and on the stalk of *Hosackia grandiflora*.

*Eudamus nevada*.—Singly or two beside each other on the leaves or stem of *Trifolium monandrum*.

*Lepisesia clarkia*.—On the underside of the leaves of *Clarkia rhomboidea* and *Gayophytum diffusum*.

*Hemaris cynoglossum*.—On the same.

*Alypia mariposa*.—On *Clarkia elegans* and *Godetia Williamsonii*.

*Alypia Ridingsii*.—On *Clarkia rhomboidea* and *Godetia Williamsonii*.

*Oncocnemis exemplaris*.—On the underside and top of the leaves of *Gentiana Newberryi* and on the grasses in the shadow of the plant.

*Anarta Kellogii*.—On the upperside of the erect, channeled leaves of *Carex filifolia*.

*Plusia californica*.—On *Trifolium monandrum*.

*Plusia Hochenwarthi*.—Mostly underneath the leaves of *Carex filifolia*, *Gentiana Newberryi*, *Salix*, *Vaccinium*, *Antennaria dioica*, *Mimulus primularioides*, *Castilleja Lemmonii* and *Pentstemon confertus*.

*Heliothis dipsaceus*.—On *Erythræa venusta*.

*Pyrausta unifascialis*.—On the stems or stalk and on the leaves. A sugar-loaf-like egg, which is laid on sideways, applied near the pointed end on *Oxytheca spergulina* and *Gayophytum diffusum*.