

wap Lake, B.C., alt. 6,000 ft.; Avalanche Mt., Selkirk Mts., B.C., alt. 7,000 ft. (*Jas. M. Macoun.*) Western Summit of North Kootanie Pass, Rocky Mts.; South of Tulameen River, B.C., alt. 6,000 ft.. (*Dr. G. M. Dawson.*) A part of the specimens referred to this variety are perhaps intermediate between it and the next.

SILENE DOUGLASII, Hook, var. *VISCIDA*, Robinson, Proc. Amer. Acad. xxvii, 145.

On slopes of high mountains at Kicking Horse Lake, Rocky Mts. (*John Macoun.*)

SILENE DOUGLASII, Hook., var. *MULTICAULIS*, Robinson.
S. multicaulis, Macoun, Cat. Can. Plants, Vol. II., p. 309.
Stump Lake, South of Kamloops, B.C. (*John Macoun.*)

SILENE GALLICA, Linn.

Common at Oak Bay, Goldstream and Victoria Arm, Vancouver Island. (*John Macoun.*) Introduced.

SILENE NOCTIFLORA, Linn.

A weed in gardens at Rupert's House, James Bay. (*Jas. M. Macoun.*) In waste places at Revelstoke, B.C., and Cedar Hill, Vancouver Island. (*John Macoun.*) Not before recorded from west of Winnipeg.

LYCHNIS AFFINIS, Vahl.

Between Lake Athabasca and Chesterfield Inlet. Lat. 63° 27', Long. 102°, 1893. (*Jas. W. Tyrrell.*) Cape Prince of Wales, Hudson Strait. (*Dr. R. Bell.*)

LYCHNIS ELATA, Wats.

Our only specimens of this species are from Avalanche Mt., Selkirk Mts., B.C., alt. 7,000 ft. (*Jas. M. Macoun.*)

LYCHNIS TAYLORÆ, Robinson, Proc. Amer. Acad. xxviii, 150.

Very slender 1 to 1½ feet high, puberulent, nearly smooth below, glandular above; stem erect, bearing 3 to 4 pairs of leaves and two or three long, slender, almost filiform 1 to 3 flowered branches; leaves thin, lance-linear, acute or