AGROTIS SUBGOTHICA.

Sir,—In reply to Mr. Tutt's note in the July number of the Can. Ent., p. 159, I would state that I have no knowledge of Haworth's work in which subgothica is described. I have everywhere taken Stephens's identifications of Haworth's species. Now Stephens figures jaculifera of Guenée, as figured in the Species Géneral, typical jaculifera, as subgothica, of Haworth (?). If, then, Stephens is wrong, and Haworth's subgothica is a variety of tritici and not our American species, this latter must be known as jaculifera, and Prof. Lintner's name of tricosa must then clearly be retained for that species, as insisted upon by myself. Stephens's figure is unmistakably based on our American species; how nearly the European tritici resembles this I cannot, at the moment, say. The following will be the synonymy of Agrotis jaculifera. According to Mr. Tutt's statement that subgothics of Haworth is a variety of tritici of Linné, our American species must be listed as follows:—

jaculifera, Guen., fig.

subgothica, Steph, fig. in err.

tricosa, Lintner.

jaculifera, Guen. var. A. jaculifera, Smith, in err.

herilis, Grote.

jaculifera, Guen. var. B.

herelis, Smith.

A. R. GROTE

SOME CORRECTIONS.

Sir,—In my paper, Can. Ent., page 152, I say I have placed Agrotis costata and its near ally A. idahoensis "together wrongly," line 10. It is clear from the context that I meant in a wrong position in my lists. The two species are closely allied, differing in colour, costata being reddish, idahoensis purplish, and, in costata, the pallid costal region is whiter and broader. The two belong together. I have always associated them, and, indeed, described the one comparatively with the other. The types are in British Museum On page 148, line 6 from bottom, for grouping, read association. On page 151, for "The Practical Entomologist," read the practical entomologists. I was writing, not of a publication, but of a class of working entomologists, whose figures (mostly duplications of the same cut) confounded the three forms: subgothica, Stephens (= jaculifera, Guen.), tricosa, Lintner (= jaculifera, Smith), and herilis, Grote (= herelis, Smith).

A. R. Grote.