Dyspepsia in Women:	FOR METRORRHAGIA:
R Ac. hydrochlorici dil m x. Ac. hydrocyanici dil m v. Liq. strychniæ m iiss. Sodii chloridi gr. xx. Glycerini m xxx.	R Ext. opii gr. jss. Tinct. hydrastis Canad
$\mathbf{A}\mathbf{q}$ \mathbf{ad} $\mathbf{\bar{5}}\mathbf{j}$.	-Lutaud, Frauenarzt; Centbl. f. d. ges. Ther.
M. ft. ht.	-
Sig.: To be taken thrice daily, an hour after meals.—Therapeutic Gazette.	FOR LICHEN URTICATUS: B. Spiritus rectificati \ Saponis mollis \ Saponis mollis \ Olei cadini \ Aquæ \ ad f\(\frac{5}{5}x\).
A LOCAL ANÆSTHETIC SOLUTIONDr. J. H.	Misce et fiat lotion.
Lowrey, of Neola, Iowa, speaks favorably of the following combination:	S.—To be applied night and morning. —The Practitioner.
B. Cocaine hydrochloride, Resorcin	Antiseptic Ointment.—Dr. P. Reclus applies R. Salol, Resorcin, Antipyrin
He has found that the use of this solution is	Boric acid 3v.

RELIABLE AND PROMPT

not followed by the systemic disturbances that

cocaine alone sometimes produces.-New York

Medical Journa!.

Two Characteristics that Commend SCOTT'S EMULSION to the Profession.

THERE ARE MORE THAN TWO but the fact that this preparation can be depended upon, and does its work promptly, covers the whole subject.

Physicians rely upon SCOTT'S EMULSION OF COD LIVER OIL WITH HYPO-PHOSPHITES to accomplish more than can possibly be obtained from plain cod liver oil. They find it to be pleasant to the taste, agreeable to the weak stomach, and rapid of assimilation. And they know that in recommending it there is no danger of the patient possessing himself of an imperfect emulsion. SCOTT'S EMULSION remains under all conditions sweet and whole some, without separation or rancidity.

FORMULA: 50% of finest Norwegian Cod Liver Oil; 6 grs. Hypophosphite of Lime; 3 grs. Hypophosphite of Soda to the fluid ounce.

SAMPLE of Scott's Emulsion delivered free to the address of any physician in regular practice.

Iodoform....

-Med. and Surg. Reporter.

Prepared by SCOTT & BOWNE, Chemists,