

produced. When the pyrexia becomes excessively high, guaiacol, used externally, lowers the temperature by inhibiting the production heat by direct absorption into the blood.

The cough, which is due mostly to nervous irritation, is greatly benefited by one-tenth to one-sixth grain of morphine sulphate, while the following inhalation, or spray, tends to materially ease both the frequency and severity of these attacks:

R.—Menthol 5 gr.
 Thymol..... 1 gr.
 Eucalyptol
 Gaultheria..... aa 10 min.
 Phenol..... 3 gr.
 White petrol. oil..... 1 oz.

Hæmoptysis has never been an alarming symptom under the administration of the following:

R.—Ac. sulph. dil..... 4 dr.
 Fl. ex. ergot..... 160 min.
 Ac. gallic..... 4 dr.
 Tr. cinnamon..... 1 oz.
 Aqua ad..... 8 oz.

Sig.—Three to four drachms every four to six hours, till hæmorrhage ceases.

This, together with the inhalation of steaming vinegar, has minimized the danger from hæmoptysis.

When there existed marked anæmia, especially seen in the so-called pre-tubercular condition, or in the early stages of tuberculosis, iron is indicated, and may usually be combined with digitalis, arsenic, and strychnia.

In dispensing with that part of this paper set apart for drug treatment, I must say that no two patients are exactly alike in any one particular, so that we treat the patient rather than the disease, and, in order to do so intelligently and conscientiously, a strict inquiry must be made into the mode of living of each case, *i.e.*, as regards exercise, food, clothing, sleeping, location and size of day and night rooms, amusements, etc. Sleeping apartments should be large, airy, having a temperature of 70° and occupied solely by the patient. It should face the east, owing to the early sunlight, and should be divested of all articles not necessary to the comfort and happiness of the patient. He should rise at 7 a.m., take exercise as ordered by the attending physician, take a light cold sponge bath, rub dry with a rough towel, and then to breakfast.

The daily food should consist of boiled or roast meats, fish, oysters soups, eggs, cod-liver oil, butter, cream, vegetables, ripe fruits (being careful of apples and bananas), liq. peptonoids. Good fresh or peptonized