Cholera, Cholera Infantum.

We have scores of testimonials, from almost every part of the world, proving the efficacy of the PAIN-KELER, when used for this terrible scourge, Cholera. If taken in time and before the disease has too firm a hold upon the system, it seldom if ever fails to effect a cure.

FOR CHOLERA.—Take a teaspoonful of the Pain-Killer in water, sweetened with loaf sugar, bathing the stomach and bowels freely with the medicine at the same time. If the attack be severe, and attended with cramps and diarrhæa, increase the dose and repeat it oftener, using also hot fomentations on the bowels as in "Cramp and Pain," &c. In the treatment of this and most other diseases of the stomach the Pain-Killer may be given with Castor Oil, in about equal proportions, with the most favorable results.

The Pan-Killer, as an internal remedy, has no equal. In cases of Summer Complaints, Dyspepsia, Dysentery, Asthma, it cures in one night, by taking it internally, and bathing with it freely. Its action is like magic.

The late Rev. Dr. Granger repeatedly expressed his belief that he owed his life to the timely use of Perry Davis' Pain-

KILLER in an attack of Cholera.

oil dd

ve

er, at.

of

ns.

es.

er. for

or

as,

en

a

d:

he oil,

ho

ng

lk

ng

es

ely

es

For Cholera Infantum.—To a child under twelve months, give from ten to twenty drops in warm milk and water, sweetened with loaf sugar, every hour, bathing the stomach in a mixture made of a teaspoonful of Pain-Killer in a half gill of milk and water, warm. If the child be between one and two years old, use for a bath equal parts of Pain-Killer milk and water, increasing the dose accordingly. For a child over two years, use the Pain-Killer clear, for a bath, increasing the dose to half a teaspoonful.