

ANESYCHIA.

A. multipunctella. N. sp.

Second joint of the palpi dark brown, tipped beneath with scattered white scales, and with a few scattered white scales above; third joint white, tipped beneath at the base with dark brown; face black; vertex white, with a central black spot; antennae fuscous; thorax white, with six black spots, one of which is placed close to the base of the wings and is continuous with the dark brown of the wings; two others on top, and one on each side a little before the tip. Primaries shining dark brown or black, with a wide white streak extending along the dorsal margin from the base nearly to the tip, and a white costal spot close to the tip; there is a sinus, or projection of the white into the dark brown portion, just before the middle, and from thence to the tip the line between the two colors is irregular, with another sinus behind the middle; there is a small black spot in the white at the first sinus, and six others along the dorso-apical margin at the base of the ciliae, and another in the costal white spot, or more properly, perhaps, there are three in the costal white spot, two of them being confluent with the dark brown color around it; ciliae white, except at the tip. *Al. ex.* $1\frac{1}{2}$ inch. Season, April and May. In nine specimens I detect no variation.

A. mirusella. N. sp.

Palpi pale yellowish ochreous; the second joint has two brown spots on its outer surface, that nearest to the base being largest; the third joint has the base and tip dark brown. Head pale yellowish or nearly white; antennae pale fuscous; thorax pale yellowish ochreous, with four brown spots, two of which are about the middle and one on each side near the tip. Primaries white, faintly tinged with ochreous yellow, and with a rather wide golden brown basal streak, which begins near the costa and diverges thence to the end of the cell, and from thence narrows and becomes more diffuse towards the apex, which it does not quite attain. The costal whitish portion is more streaked and suffused with ochreous than the dorsal portion, which has a small brown spot before the middle; there is likewise a row of small brown spots around the apex, at the base of the ciliae. *Al. ex.* $1\frac{1}{2}$ inch. Season, April and May.

I have not examined the neuration, and the ornamentation is unusual in the genus; but the other characters are those of *Anesychia*.