

*Bradynotes montanus*, n. sp.

Small, comparatively slender, dark reddish brown, with livid yellow and white markings; tegmina sometimes obsolete, when present as in *Pezotettix gracilis*.

Vertex between the eyes very broad, slightly deflected, scarcely sulcate, with a very faint though perceptible transverse depressed line joining the upper extremities of the eyes, this line bending forward so as to form a gentle arc with the convexity to the front. Frontal costa moderate, nearly equal throughout, expanding a little at the ocellus, where it is very shallowly sulcate. Pronotum simple, expanding equally and rapidly posteriorly in the female, nearly equal in the male; front margin straight, posterior margin slightly but broadly truncate; median carina of the pronotum slight, visible throughout, on the anterior lobe as a mere smooth line not elevated in the least, in some specimens represented by a very faint depressed hair line, on the posterior lobe slightly elevated, the sides gently tapering; lateral carinae obsolete, except in some specimens where they are visible on the extreme front edge of the anterior lobe as blunt shoulders. Anterior lobe coarsely and somewhat distantly, and the posterior minutely and thickly punctate, rugulose. Abdomen carinate, tapering gently and evenly backward. Tegmina situated low on the sides, small, straight, narrow, three times as long as broad, the apex rounded. Posterior femora short, somewhat tumid, nearly (female), or just reaching the tip of the abdomen (male). Last ventral segment of the male abdomen upturned, prow-shaped, entire; cerci elongate conical, with the tips directed backward and gently downward, reaching a trifle beyond the tip of the supra-anal plate. This latter very similar to that of *B. obesus*, but proportionally broader apically. Female cerci mere rudiments, while in *B. obesus* and *B. opimus* they nearly equal those of the male.

The general color is dark reddish brown with livid yellow and white markings above, flavous beneath. Face griseous yellow, becoming darker above, the vertex and occiput brownish fuscous; a very narrow but sharply defined yellow line commencing near the upper posterior angle of the eye, separating the occiput from the genae and extending backward on to the pronotum, where the lateral carinae would be if present; in some specimens a third line of a similar nature is present, beginning at the vertex and extending backward along the middle of the occiput. Sides of pronotum yellowish white, with a broad brownish piceous band extending from near the middle of the anterior lobe obliquely upward to the posterior