

*Andrena pluvialis*, Ckll., CAN. ENT., XXXIII, 1901, p. 154.

♀, 1st May, 1894, Olympia, Wash. (T. Kincaid); Victoria, B. C. (G. W. Taylor); 29th May, 14th June, 1896, Livingston, Vanc.

*Andrena transnigra*, n. sp.

♀ 11 mm.

Type locality: Seattle, Washington. Type Am. Ent. Soc., Phila.

♀, 17th April, 1896, the type.

*Andrena Seattlensis*, n. sp.

♀ 10 mm.

Type locality: Seattle, Washington. Type Am. Ent. Soc., Phila.

♀, Seattle, Wash. (T. Kincaid); 13th June, 1902, Vancouver, B. C. (Harvey); 17th May, 16th June, 1896, Livingston, Vancouver, Victoria, B. C. (G. W. Taylor); 20th May, 1895, Almota, Wash. (C. V. Piper).

*Andrena Chapmanæ*, n. sp.

♀ 12 mm. Pubescence black. This species is included to show the relation between it and the form from Oregon, which may be a distinct race.

Type locality: Yosemite, California. Type Acad. Nat. Sci., Phila.

24th June, 1902 (B. Chapman).

*Andrena Chapmanæ* race.

1st June, 1897, Corvallis, Oregon (Cordley).

*Andrena Pullmani*, n. sp.

♀ 10 mm. Face with black dorsum with white pubescence.

Type locality: Pullman, Washington. Type Univ. Nebraska.

♀, 14th April, 1897; 6th May, 1898 (C. V. Piper); ♂, 25th April, Livingston, Vanc.; 3rd May, 1896, Elkton, Or.

*Andrena longihirtiscopa*, n. sp.

♀ 10 mm.

Type locality: Vancouver Island. Type Acad. Nat. Sci., Phila.

♀, Vancouver Is. (G. W. Taylor).

*Andrena vicinoides*, n. sp.

♀ 12 mm.

Type locality: Victoria, B. C. Type Acad. Nat. Sci., Phila.

♀, Victoria, B. C. (G. W. Taylor); 2nd June, 1897, Olympia, Wash. (Kincaid).