ception in Nymphæa, Nuphar, Butomææ, Lardizabaleæ, ibid., 555; Hydropeltideæ, Mesembryanthemum, 556; explanation of, 556; belong to the transformed leaf, or carpel, and not derived from processes of the axis, 563; arrangement of in Cyrtandraceæ, 558; in Hydnora, Rafflesia, and Brugmansia, ibid.

- Oxalideæ, distinction of the order, observation on its affinities, and on the genera composing it, 113-14
- Palmæ, observations on the order and on its distribution in Terra Australis and New Zealand, 51; further observations on the order, on its distribution, and on the species found in the vicinity of the Congo, 14C
- Paniceæ, a tribe of Gramineæ, observations on the tribe and its distribution in Terra Australis and elsewhere, 57-8
- Papaveraceæ, stigmata and placentæ of adjoining cells confluent in, 559
- Papaw of the banks of the Congo, probably brought from America, 155; arguments in favour of its American origin, 158
- Papilionaceæ, observations on the order, and on its distribution in Terra Australis, 24; species found in the vicinity of the Congo, 111; in Central Africa, 295
- Parasites on roots, distinguishing characters of, in their leaves, seeds, and embryo, 391-2; modes of union between the stock and parasite, 392; no instance, at present, of parasites on roots originating from other parts of the plant, 395; found, however, to be the case in Rafflesia Arnoldi, 403.
- Park, Mungo, plants collected by him on the banks of the Gambia, 152

- Parkia, characters of the genus and observations on the species composing it, 289 and note
- Parnassia, atigmata and placentæ of adjoining cells confluent in, 559
- Paropsia, its affinities, 121
- Parry, Captain, list of plants collected at Melville Island, during the voyage of, 183, 195.
- Passifloreæ, observations on, 121; regarded as forming a natural class, with Cucurbitaceæ and Homalinæ, 122; observations on the African species of, 150; compared with Rafflesia, 386-9
- Paterson, Colonel, plants collected by him in New South Wales, and at Port Dalrymple, 6
- Peliosanthes Teta, its seed becomes naked by the early rupture of the pericarpium, 362
- Pennisetum trichotomum, a great annoyance to man and beast, 302-3
- Petaloma belongs to Melastomaceæ, 118
- Phænogamous plants, number of species in Persoon's Synopsis, 5; number of Australian species known [in 1814], 7; [in 1849], 338
- Philydrum, observations on the genus, 53
- Phyllocladus, analogy of male and female organs in, 460
- Phytolaceæ, establishment of the order, observations on its structure and affinities, and on the species found in the vicinity of the Congo, 138
- Pineapple of the banks of the Congo brought from America, 155
- Pinus, origin of the ovulum in, 458; analogy of male and female organs in, 461; plurality of embryos in several species of, 568-572; the species of, well adapted for an investigation into the origin and development of the embryo, 569; early changes consequent on impregnation in, 569-571; separation of the amnios, 570; formation of cavity in, ibid.; development of funiculi, 577; formation of embryos, ibid.; ramification of funiculi, 577 8; each impregnated ovulum con-