pseuderminia Peck.

1823-Peck., Mass. Agr. Rep. & Journ., VII., 328, pl. 1, Arctia.

1837—Westw., ed. Drury, I., 7, pr. syn.

1855—Wlk., C. B. Mus., Lep. Het., III., 667, Spilosoma. caprotina Drury.

1770-Dru., Exot., I., pl. 3, f. 3, Phaiæna.

1782—Cram., Pap. Exot., III., 170, pl. 287, f. C, Phalana.

1797—Sm. & Abb., Ins. Ga., II., 133, pr. syn.

1837—Westw., ed. Dru., I., 7, pr. syn.

californica Pack.

1864-Pack., Proc. Ent. Soc., Phil., III., 125, Leucarctia.

1872—Pack., 4th Rept. Peab. Ac., 86, pr. syn.

packardii Schaupp.

1882-Schaupp, in Check List Bkln. Ent. Soc., 8, Spilosoma.

Habitat-United States and Canada.

Food plant-Omnivorous.

## L. albida Stretch.

1873-Stretch, Zyg. & Bomb., 203, pl. 8, f. 22, Leucarctia.

1875—Butler, Cist. Ent., II., 36, an sp. dist. virginica?

1876—Stretch, Rept. Geol. Surv., west 100 merid., V., 798, pl 40, ff. 4 and 5, Leucarctia.

Habitat-California.

The Arctia scuirus of Boisduval was referred by Grote and Robinson to acræa, but is now recognized as referring to Euchaetes collaris! The peculiar male characters of the acræa have been described by Mr. Morrison, and later figured by myself.

Leucarctia permaculata Pack. is an Ecpantheria, and is referred to under that genus. I do not find the species in Mr. Grote's list, but perhaps it may be referred to in some paper which I have not seen.

(To be continued.)

Mailed July 4th.