

repeated; fringes brown. Body rusty brown. Tooth on internal margin of primaries not prominent. Expanse 36 m. m. Collection Buff. Soc. Nat. Sci.

This seems to be a shorter and broader-winged form than the European, in which it would conform to Dr. Speyer's law of variation in the Nocturæ. The outer line is less distinct than in the European species, of which it may be a modification.

### ON PSEUDOHASIS HERA (HARRIS).

BY A. R. CROTE, BUFFALO, N. Y.

Through the kindness of Lieut. Carpenter, I have received from Yellowstone Park a ♂ specimen which agrees with Mr. Walker's description of *pica* (B. M. Lists, 1318), and must be the same, although outside of the common black outer fascia the wings are tinted with yellow, while appearing white at first glance. But Walker's description seems identical with Harris' *Hera*, and his specimen is from Doubleday also. Probably the same specimen has been made the type of both species. Walker's locality, "United States," is indefinite, and probably not as originally given by Doubleday. Further comparisons are needed to settle the differences between *eglanterina* and *Hera*, which I have considered (Am. Phil. Soc., 1874, p. 4) as specifically identical.

It is not necessary, at this time, to analyse Audubon's plate; for the two names, *Hera* and *pica*, are founded on *specimens*, and no name is attached by Audubon or others to the original figures. I would correct the synonymy of the two forms as previously given by me.

*Pseudohasis* G. & R. (1866).

(Type: *Saturnia eglanterina* Boisd.)

1. *Pseudohasis Hera* (Harris) G. & R., Ann. Lyc. N. Hist., N. Y., 8, 377; *Saturnia Hera*, Rep. Ins. Mass., 286, 1841; *Hemiluca pica* Walk., B. M. Lists, 6, 138.

Hab. Rocky Mountain Region.

2. *Pseudohasis eglanterina* (Boisd.) G. & R., Ann. Lyc. N. Hist., N. Y., 8, 377; *Saturnia eglanterina* Boisd., Ann. Soc. Ent. Fr., 2nd Ser., x, 323; *Telca eglanterina* H.-S., Exot. 60, fig. 445.

Hab. California.