be now very accurately measured-in length, as I said before, 134 inches, and 24 inches broad, extending from the pubis to the first false rib in an oblique direction. passing backwards. The wound in the scrotum still open, but the communication between the abscess and the extreme opening seems closed.

25th .- Opened the abscess this day, at the pubis. About 3 viij of very fætid, greenish matter escaped; it came away

with a gurgling sound; wound in the scrotum progressing favorably.

27th .- Abscess still discharging; ordered beer, and a generous diet.

Sept. 2d .- The opening of the abscess has closed; the wound in the scrotum nearly healed.

8th .- Has resumed work, being quit well.

St. Martin, Isle Jesus, Sept. 11, 1851.

ART. XXI. -- Case of Elephantiasis Arabica of the Penis and Scrotum, by S. C. SEWELL, M.D., Edin. L.R. C.S.E., Lecturer on Clinical Medicine, University of McGill College, &c., &c.

Pierre Houlée, æt. 23, laborer, presented himself at the Montreal General Hosnital, on the 22nd March, 1851, with elephantiasis arabica of the penis and Fig. 1, is a very accurate representation of his appearance, taken previous to the operation by Mr. Bucke, medical student, and transferred to wood by Mr. Walker. The scrotum was the size of a new-born infant's head; the penis

a-half inches, measured fourteen-and-ahalf inches in circumference at the junction with the pubis (1), eight-and-a-half inches at the first notch, which corresponded with the corona glandis; it then increased somewhat bulgingly, and from (2) to extremity (3) it measured uniformly nine-and-a-half inches, terminating abruptly as if truncated. Over the left half of the body of the penis, the skin was sound; the right half was transformed into a tumor, which, like the scrotum, presented a surface covered with large flat The prepuce, which was tubercles. elongated nine-and-a-half inches, conwas elongated, to the extent of twelve-and- sisted of pedicled tubercles on a thickened

