

pag. lin.

- an annual, is very variable in its size, colour, and number of florets in the ray. Native specimens collected by Mr. Bradbury are covered with rigid hairs, and have but one flower. Cultivated plants raised from seeds of the same species are more smooth, branch more or less; the leaves are deeper divided, and produce several flowers double the size.
- 575 *Rudbeckia columnaris*, adde *R. columnaris*; foliis pinnatifidis: foliolis lanceolatis, exterioribus ordinatim majoribus, receptaculo cylindraceo elongato. *Sims in Bot. mag.* 1601. This species is not above a foot high, and appears to be annual.
- 580 *Iva ciliata*, adde *Iva annua*. *Mich. fl. amer.* 2. p. 184.
- 585 *Orchis ciliaris*, adde *Habenaria ciliaris*. *Brown in Ait. kew. ed. 2. vol. 5. p. 194.*
- Orchis cristata*, adde *Habenaria cristata*. *Brown in Ait. kew. ed. 2. vol. 5. p. 194.*
- 587 *Orchis viridis*, adde *Habenaria viridis*. *Brown in Ait. kew. ed. 2. vol. 5. p. 192.*
- Orchis bracteata*, adde *Habenaria bracteata*. *Brown in Ait. kew. ed. 2. vol. 5. p. 192.*
- 588 *Orchis hyperborea*, adde *Habenaria hyperborea*. *Brown in Ait. kew. ed. 2. vol. 5. p. 193.*
- Orchis fimbriata*, adde *Habenaria fimbriata*. *Brown in Ait. kew. ed. 2. vol. 5. p. 193.*
- 589 *Neottia cernua*, adde *Bot. mag.* 1568.
- Neottia repens*, adde *Goodyera repens*. *Brown in Ait. kew. ed. 2. vol. 5. p. 198.*
- 590 *Neottia pubescens*, adde *Goodyera pubescens*. *Brown in Ait. kew. ed. 2. vol. 5. p. 198.*
- Arethusa divaricata*, adde *Pogonia divaricata*. *Brown in Ait. kew. ed. 2. vol. 5. p. 203.*
- 592 *Cymbidium pulchellum*, adde *Calopogon pulchellus*. *Brown in Ait. kew. ed. 2. vol. 5. p. 204.*
- Cymbidium verecundum*, adde *Bletia verecunda*. *Brown in Ait. kew. ed. 2. vol. 5. p. 206. Bot. mag.* 930.
- 594 *Cypripedium parviflorum*, adde *Bot. mag.* 911.
- 595 *Cypripedium arietinum*, adde *Bot. mag.* 1569