

**Tumor of the Middle Lobe of the Cerebellum.**—Dr. MEYERS presented to the Society a young man where the above diagnosis had been made. (See page 35.)

Dr. GRAHAM said that from what he had learned of the case as presented, he would lean to the diagnosis of ataxic paraplegia. The papillitis and headache might have been due to meningitis.

Dr. ATHERTON asked an explanation of the suddenness of the attack.

Dr. CASSIDY thought the diagnosis of the tumor, as tubercular, was negated by the absence of tubercle in the rest of the body.

Dr. ROSS described a case he had seen of tubercular tumor of the middle lobe of the cerebellum, to which the case reported bore some resemblance. However, he leaned to the diagnosis of ataxic paraplegia in this case.

Dr. MEYERS pointed out that his diagnosis rested mainly on the optic atrophy, the marked decrease in the number of respirations, the limited ataxia, the strength of the leg muscles. The original trouble might have been peripheral neuritis.

**Sarcoma of the Ovary.**—Dr. ROSS presented a specimen of ovarian tumor from a negress. She had suffered from it five years, when suddenly she experienced a rise in temperature of  $105^{\circ}$ , and great abdominal pain. It looked perfectly healthy on removal, but upon opening it was seen to be undergoing sarcomatous degeneration.

---

## Toronto Medical Society.

(JANUARY 17TH, 1895.)

*President, DR. PETERS, in the chair.*

**Scleroderma.**—Dr. MCPHEDRAN presented a patient suffering from scleroderma. It commenced about nine months ago, a white spot being first noticed in the forehead, which presented something of the appearance of a chalk mark. It extended upward upon the head about two inches within the margin of the hair line. Its width was about half an inch. The patch became completely bald. Thickening of the skin and the tissue has taken place. It is slightly anæsthetic.

**Urethral Calculus.**—Dr. PETERS showed a urethral calculus which he had removed from a boy seven years of age. The history of its presence lasted from the patient's birth. At the age of four he received a blow in the perineum. Since then he has had frequent passages of bloody urine accompanied by great pain. The urine kept continually running away. On passing a sound the stone could be distinctly felt