

# The Canadian Entomologist.

VOL. XVI. LONDON, ONT., FEBRUARY, 1884.

No. 2

## NOTES ON SPHINGIDÆ CAPTURED AT ORONO, MAINE, AND VICINITY.

BY MRS. C. H. FERNALD.

- Lepisesia flavofasciata*, Barnst.—Very rare, taken in the hottest part of the day, on lilac and apple blossoms; also on *Amelanchier Canadensis*.
- Hemaris diffinis*, Boisd.—Common on Tartarian honeysuckle and lilac, in the middle of the day.
- Hemaris gracilis*, G. & R.—Quite rare, found on the same flowers and at the same time as *H. diffinis*, and also flies about sunset.
- Hemaris thysbe*, Fabr.—Very common, flying in company with *H. diffinis*.
- Amphion nessus*, Cram.—Quite common in company with the last named species.
- Deilephila chamænerii*, Harris.—Some seasons quite common on Tartarian honeysuckle and lilac, at twilight.
- Deilephila lineata*, Fabr.—Taken on *Oenothera Lamarckiana* just after sunset, in September.
- Philampelus pandorus*, Hübn.—One example bred from larva on grape vine.
- Philampelus achemon*, Drury.—Larvæ found on grape vine, but failed to emerge.
- Everyx chærilus*, Cram.—Rare, at sugar, in July.
- Smerinthus geminatus*, Say.—Quite common at light, in June.
- Smerinthus cerisyi*, Kirby.—Very rare, at light, last of May.
- Paonias excæcatus*, A. & S.—Quite rare, at light, in June.
- Calasymbolus myops*, A. & S.—Two examples taken at light.
- Triptogon modesta*, Harr.—Rare, at light, first of July.
- Ceratonia amyntor*, Hübn.—Quite rare, on climbing honeysuckle.
- Daremma undulosa*, Walk.—Quite common, at light and on flowers in June.
- Phlegethontius cingulata*, Fabr.—Two examples on flowers of *Nicotiana affinis*, quite late in the evening, in the first of Sept.