

lesser extent beyond the median band in the lower half of the wing. A faint ochraceous curved band leaves costa, above orbicular, passing downward and inward to base, almost at inner margin, dividing this portion of wing in about two equal parts, the lower half being somewhat more tinged with ochraceous than the upper. Fringe white, with ochraceous blotches between veins. Hind wings yellowish-white, more or less tinged with fuscous, especially outwardly. Fringe white.

Beneath, fore wing blackish centrally, paler along costa and outer margin. Hind wings pale yellowish-white.

Collar, patagia and thorax creamy-white, with ochraceous shading. Thorax with posterior tuftings. Abdomen fuscous-yellow.

Type, one ♀, Southern New Mexico, from Mr. Poling.

*Chamaeclea gladiola*, n. sp.—Expanse, 28 mm.

Fore wing, at base, beyond s. t. line and a large oval patch in the centre below costa, creamy-white. Remainder of wing dark olivaceous-brown, with an admixture of violet and paler olivaceous and yellow scales. Ordinary lines not distinctly marked. Basal portion of wing is creamy-white except along costa, where it is of the same shade as the median portion of wing. The reniform is present at outer edge of oval white patch, though not very plainly marked. It is pale-ringed, with dark centre, narrow and upright. Indications of a row of terminal black intravenular dots. Fringe white. Hind wings blackish-brown, with faint indications of mesial band and discal dot. Fringe white.

Beneath, fore wings blackish, with central yellowish patch, yellowish along costa. Hind wings yellowish-white, with dusky mesial band. Collar yellowish at base, remainder of collar, patagia and thorax white. Thorax posteriorly with some olivaceous and violet scales.

Types, ♂ and ♀, Santa Catalina Mts., Ariz.

This species should stand next to *Anthæcia scira*, Druce, Biol. Centr. Amer. Het., Pl. 28, fig. 5. The type of maculation and colours are about the same, but the Mexican species entirely lacks the large white patch in centre of wing.

*Oxycnemis acuna*, n. sp.—Expanse, 15 mm.

Ground colour gray, more or less covered with brownish scales, markings black. Ornamentation similar to other species of the genus. Claviform long linear, gray, narrowly outlined in black, brownish centered. Orbicular similar in form and size to claviform, gray, narrowly outlined in black, brown centered, lying parallel to and extending a little beyond claviform. Reniform gray, with brown centre, surrounded by a few black