



BAYER'S PHARMACEUTICAL PRODUCTS

SOMATOSE A tasteless, odourless

(Trade Mark.) nutrient meat powder; it contains all the albuminoid principles of the meat in an easily soluble form. It has been extensively employed and found to be of the greatest service in Consumption, diseases of the stomach and intestinal tract, Chlorosis and Rickets. It is of great value in convalescence from all diseases. **SOMATOSE** strengthens the muscles and stimulates the appetite in a remarkable manner. **SOMATOSE** has been found to act as a most efficient galactagogue. Dose for adults: a level teaspoonful three to four times a day with milk, gruel, coffee, etc.

IRON SOMATOSE (Ferro-Somatose). A first-class tonic, containing the albuminous substances of the meat (albumoses) organically combined with iron. Special indications: Chlorosis and Anæmia. Daily dose: 75 to 150 grains.

MILK SOMATOSE (Lacto-Somatose). A strength-giving food containing the albuminous matter (albumoses) of the milk. Daily doses for children: 1 to 2 teaspoonfuls; for adults: 2 to 3 tablespoonfuls.

TRIONAL (Diethylsulfonmethylethylmethan). A most reliable and quickly-acting hypnotic of the Sulfonal group. Dose: 16 to 20 grains, in a large cup of hot liquid.

IODOTHYRINE The active principle of the thyroid gland. It is most efficacious in Strumous Diseases, Myxœdema, Obesity, Rickets, Psoriasis, Eczema, and Uterine Hæmorrhages. Dose: 5 grains two to eight times a day for adults; 5 grains one to three times daily for children.

LYCETOL (Tartrate of Di-Methyl-Piperazine). Anti-Arthritic, Uric Solvent. Has a marked effect on the diuresis. Dose: 16 to 32 grains daily.

ARISTOL (Dythyroidiodide). A Cicatrissant which is an excellent, odourless substitute for iodoform and highly recommended for Burns, Wounds, Scrofulous Ulcerations, etc.

EUROPHEN (Isobutylorthocresoliodide). A perfect substitute for Iodoform. Odourless and non-toxic. Has a covering power five times greater than iodoform. Especially useful in Ulcus molle et durum.

PROTARGOL A new silver preparation. Most reliable in cases of Gonorrhœa. Antiseptic wound healer. Excellent results in cases of Gonorrhœal Ophthalmia. Solutions of $\frac{1}{4}$ to 2 J Ointments.

LOSOPHAN (Triiodometacresol). Particularly efficacious in the treatment of all kinds of cutaneous disorders caused by animal parasites.

TANNIGEN (Triacetyl of Tannin). An almost tasteless intestinal astringent. Most efficacious in Chronic, Acute and Summer Diarrhœas. Adult dose: 8 grains every three hours.

TANNOPINE (A new intestinal astringent). (Formerly "Tannone"). Special indications: Tuberculous and non-tuberculous Enteritis, Typhus. Dose: 15 grains, three or four times daily.

SALOPHEN (Acetyl of Para-Amidosalol). Specific for Influenza, Headache, Migraine, Acute Articular Rheumatism, Chorea, Sciatica. Dose: 15 grains, four to six times daily. In powders, etc.

ANALGEN (Ortho-Ethoxy-ana-Monobenzoylamidoquinoline). A specific for Malaria. Highly recommended in Acute Rheumatism of the Muscles, Sciatica, Facial Neuralgia, etc. Malaria: before the paroxysm of fever 20 to 30 grains; between the fevers 15 grains every 3 hours. Rheumatic affection and Sciatica: 15 grains, 4 to 5 times daily. The use of **ANALGEN** is accompanied by a reddish coloration of the urine, which, however, is not produced by the presence of blood corpuscles. The red color of the urine may be avoided by taking alkaline waters.

PHENACETINE-BAYER (Acetyl of Para-Phenetidin).

PIPERAZINE-BAYER (Diethylene-diamine).

HEROIN (Di-acetic ester of morphine). An excellent substitute for codeine. In doses of 0.005 gramme, 3 to 4 times daily, it has given excellent results in cases of Bronchitis, Pharyngitis, Laryngitis, Catarrh of the Lungs in phthisical persons, and in Asthma Bronchiale. In the latter two cases, the dose may be increased to 0.01 gramme.

CREOSOTAL (Creosotum carbonas puriss). A mixture of the phenol carbonates of creosote. Most valuable in tuberculosis of the lungs. Doses of $\frac{1}{4}$ to 5 drachms per day, in wine, brandy, or cod liver oil.

DUOTAL (Guaiacolum carbonas puriss). Great success in cases of Pulmonary Phthisis. Doses of 8 to 96 grains per day.

SULFONAL-BAYER (Diethylsulfondimethylmethan).

SALOL-BAYER (Phenyl Ether of Salicylic Acid).

Samples and literature may be had on application to the

DOMINION DYEWOOD & CHEMICAL CO., TORONTO.

Sole Agency and Depot in Canada for all "BAYER'S" Pharmaceutical Products. (Wholesale only.)