

medium albo-pubescente, apice truncatis, et 3 vel 4-denticulatis; pedibus picis, antennis flavo-testaceis. Long. 4.5 mm.

Lower California, Mr. Ulke. A singular species, but resembling entirely in form and size *G. minuta*. The elytra are distinctly truncate at tip, and the truncature is serrate, the suture and outer angle being prominent, with one or two intermediate cusps.

CALLIDIUM FABR.

485. *C. vile*. Nigrum, subtiliter cinereo-pubescent, prothorace latitudine vix brevior, lateribus rotundatis, fortiter punctato; elytris sat dense fortiter punctatis, et transversim rugosis; antennarum articulo 2do sequente triplo brevior. Long. 4.5 mm.

Mendocino, California; collected by Mr. Behrens. Readily known by the small size, black color, and coarse sculpture. The thighs are strongly clubbed, the front coxæ are contiguous, and the mesosternum obtusely triangular. I cannot see the mesonotum, but I have no doubt from the other characters that there is no stridulating surface.

486. *C. hirtellum*. Elongatum minus depressum, nigrum nitidum, pubescens, et pilis nigris erectis villosum, prothorace ferrugineo, punctato, callis lævibus haud elevatis ornato, latitudine paulo brevior, lateribus rotundatis; elytris conoide punctatis sæpe testaceis; antennis pilosis, basi rufescentibus, articulo 2do 3io dimidio brevior. Long. 8 mm.

Nevada, Mr. Ulke. Two specimens, one of which is black, with the prothorax ferruginous; the other has also the elytra brownish ferruginous. The mesonotum is smooth and polished, with a few scattered punctures; the mesosternum is triangular; the thighs strongly clubbed, the front coxæ contiguous. *C. æreum* Newman (*pallipes* Hald.), belongs to the same division of the genus, characterized by the rather stout antennæ, not thickened but very hairy towards the base, with the 2d joint about half as long as the 3d, and the punctuation less dense than in the metallic blue species.

XYLOCRIUS LEC.

487. *X. cribratus*. Ater, pube nigra villosus, prothorace fortiter punctato, latitudine brevior, postice subangustato, lateribus antice valde rotundatis, postice subsinatis; elytris subreticulatim grosse punctatis, punctis versum apicem paulo minoribus. Long. 12.5 mm.