

South Louisiana," *Psyche*, vol. iv., p. 219-222). For the identification of the species (except those marked with an asterisk) I am indebted to Dr. G. H. Horn, who kindly examined them for me.

CICINDELIDÆ.

Tetracha carolina (Linn.) One full grown larva dug from its hole, 10th April. Milnburg Road.*

Cicindela repanda Dej. One specimen seen (not taken) in an unpaved side street in New Orleans, first part of April. Same species seen also at the mouth of the Red River in Louisiana, 22nd June, and taken at Vicksburg, Miss., 23rd June.

" *tortuosa* Dej. One specimen taken on a low, sandy mud-flat of the Mississippi just below the mouth of the Red River in Louisiana, 22nd June. In company with *repanda*. This is the only species in the list which was not taken (or seen) on or south of parallel 30°.

CARABIDÆ.

Scarites subsriatus Hald. Of occasional occurrence under logs in moist places, N. O.

" *subterraneus* Fabr. Same with preceding, N. O.

Brachynus lateralis Dej. A number taken under old wood on the edge of a wet forest, B. la F.

" *tormentarius* Lec. In colonies, often in company with *Galerita*, under logs (old railroad ties) and pieces of wood in low, moist places, on the edge of the swamp especially. †

" *fumans* (Fab.) A number taken under logs in drier places than preceding. N. O.

Panagæus cruigerus Say. One specimen under a log on the edge of a muddy, wet forest, 4th May. B. la F.

Galerita janus Fabr. This and the next species in same places with *Brachynus tormentarius*; often found in company with that species.‡ Also overtaken running through passages made in a

* There are dummy roads running through the swamp from New Orleans to Milnburg, Spanish Fort and West End, three resorts on Lake Pontchartrain.

† Where no locality is given, the species will be understood to have been taken both in vicinity of New Orleans and on Bayou la Fourche.

‡ Where no date is given, unless otherwise stated, the species was more or less common during the time of my stay.