A few specimens in the Museum of the Canadian Survey, which Dall thinks are "probably the true *V. sincera* of Say, or a variety of it," were collected on the island of Anticosti by Professor Macoun in 1884, and on the Attawapiskat and Kawinogans rivers by Mr. McInnes in 1904. Those from the Kawinogans River have the outer half of the last volution free and partially uncoiled.

Limnæa stagnalis appressa Say.

Stewart River, near Mayo River; two specimens.

Limnwa Randolphii Baker.

Frances Lake, eleven fine and large specimens; Finlayson Lake, eleven specimens, mostly immature; Lewes River, one small specimen; and Lake Marsh, Lewes River, seventeen fine and mostly adult shells.

Limnæa palustris Muller.

Frances Lake, two specimens.

Limnæa Vahlii Beck (Dall).

Frances Lane, nine specimens; Finlayson Lake, twelve specimens.

Limnæa arctica Lea (Dall).

= Limnæa Pingelii Beck, var. (Dall).

Stewart River, near Mayo River, two specimens.

Planorbis trivolvis Say.

Stewart River, near Mayo River, seven specimens of a rather large, depressed and thin-shelled form of this species, with the spiral angulation obsolete.

Segmentina armigera Say.

Stewart River, near Mayo River; one specimen.

Physa sp. indet.

Finlayson Lake; one very immature specimen.

Ottawa, April 5, 1905.