Driotura gammeroidea, var. fulva, n. var.—Size and form of the species larger than var. flava. Entirely brownish fulvous, except the eyes, which are darker.

Described from eight specimens from Denver, Colo., collected by the author.

Driotura robusta, var. vittata, n. var.—Size and form of the species, black and white, variable. Vertex with a transverse light line on anterior margin, expanded into two spots at apex; four oblique black stripes on elytra, alternating with four light ones. A transverse light band on abdomen, and a broader one on face.

Described from six examples from Southern Colorado.

Acinopterus acuminatus, var. variegatus, n. var.—Form and structure of the species, but much lighter coloured. Vertex, pronotum and scutellum inclined to be reddish, especially in the male. Elytra whitish pruinose, nervures greenish, not margined, except towards apex and along the sutural margin, three fuscous points along the suture, and sometimes one on the disc of each elytron.

. Described from twenty-four specimens from Colorado and Arizona.

A. acuminatus, var. viridis, n. var.—Form and structure of the preceding nearly; slightly smaller. Bright grass-green both above and below. Eyes and extreme tip of elytra fuscous.

Described from a number of specimens from Southern Colorado and Arizona. This is the common form in Southern Colorado, where it was collected by E. P. Van Duzee and the author.

A. acuminatus, var. brunneus, n. var.—Slightly larger than the preceding variety. Vertex, pronotum and scutellum pale green, washed with cinnamon-brown. Elytra pale cinnamon-brown, slightly fuscous at tip. Whole insect with a slight tawny iridescence, below pale green.

Described from three specimens from Rifle, Colo.; taken by the author.

Liburnia Slossoni, n. sp.—Resembling Stenocranus lautus in size and general appearance. Somewhat resembling D. maidis. Length, macropterous 2,5 mm. Face broad, strongly carinate, slightly narrowing above. Elytra very long and narrow, resembling a Stenocranus, the outer branch of the first and the inner branch of the third sector uniting with the cross nervure alongside the second sector. Colour: Face black, the carinæ light, basal compartment of vertex, pronotum and