

Abdomen ovate, polished, short petiolate, black, except the greater part of the first segment and the extreme tip, which are black; second, third and fourth segments sub-equal in length. Legs rufous, the tarsi lighter; femora, especially the anterior pair, very stout; anterior tarsi chelate, first joint long; anterior femora with a black mark below at the base, tips of posterior femora and tibiæ also black. Wings hyaline, with a fuscous band just beyond the stigma, also a faint fuscous spot at the apex of the second basal cell, stigma lanceolate.

Described from a single female specimen collected at Mosholu, N.Y., July 25, 1903, by Mr. J. R. de la Torre Bueno.

This species may be distinguished from the only other described species of *Bocchus* (*B. flavicollis*, Ashm.) by its colour, larger size, different length of abdominal segments and configuration of the antennæ. It agrees very well with the generic diagnosis given by Ashmead (Monog. Proctotrypidæ, p. 91).

*Oxylabis bifoveolatus*, sp. nov.—Male. Length, 3 mm. Black; legs fuscous, varied with rufous. Head shining black, rugulose on the occiput, with a short impressed longitudinal line above the ocelli; head margined behind; cheeks shagreened; mandibles dark fuscous. Antennæ 14-jointed, distinctly longer than the body, tapering toward the tips; two basal joints black, the rest fuscous. Scape very short, twice as long as the pedicel, which is less than one-half as long as the first flagellar joint; first eleven flagellar joints of equal length, the last one and one-half times as long as the preceding and more slender. Sides of the pronotum coarsely and obliquely striate. Mesonotum with two deep furrows, less pronounced anteriorly, and with a triangular fovea just before the scutellum. Scutellum with two deep foveæ. Post-scutellum with a median groove and a more delicate one on each side; its tip produced into a long, acute black spine, which is perpendicular to the posterior face of the metathorax. Metanotum anteriorly coarsely rugose-striate, at the middle with a widely-interrupted transverse furrow, behind this with a fovea on each side, and medially at the tip with a large enclosed space. Petiole on abdomen suddenly constricted in front, a little wider than long and coarsely fluted. Remainder of abdomen compressed and very shining, impunctate; second segment very long, others short. Legs fuscous; front tibiæ, knees and bases of the tarsi lighter. Wings