

more so. Soon after the operation the patient suddenly developed a cough with expectoration, which soon became purulent, and was at times bloody. Three weeks after the operation a large rubber drainage-tube was passed through the Douglas pouch and out by the vagina, being carried a few inches beyond the vulva. This did no good. She gradually sank, and died exhausted six weeks after the operation. Two days before death she complained of sore throat, and on examination the fauces, tonsils and posterior wall of the pharynx were found to be covered with a diphtheritic membrane. Until the autopsy, I adhered to the original diagnosis of suppurating, universally adherent, ovarian cyst. Dr. R. J. B. Howard, acting pathologist to the Hospital, made the autopsy. I append his report:—"On opening abdomen a large globular mass presents, of the size of a man's head, occupying false pelvis; this and the parietes are everywhere covered by a grey, rough membrane about one-eighth of an inch thick. The transverse colon is firmly adherent to the upper surface, and is also bound tightly down to the liver. A collection of pus is found below and by the side of the spleen, and another, smaller, under left lobe of liver in middle line. The anterior peritoneal cavity is thus converted into a suppurating cyst, extending from liver down into true pelvis, nearly filled by the mass, which is found to consist of all the intestines, except the transverse colon, closely matted together by recent slight adhesions, which are studded with miliary tubercles. The cyst wall is apparently much older than the inter-intestinal adhesions, and looks like an unhealthy granulating membrane. The walls and viscera of true pelvis are covered by the same membrane. The great omentum has quite disappeared; but no doubt had been spread out over the intestines, and formed part of the membrane covering them. All the abdominal viscera adherent to one another and to parietes. Liver fatty; contains a few gray granulations. Kidneys contain a few gray granulations. Lungs universally adherent; abundantly studded with gray granulations. Tonsils and pharynx—surface gray and sloughy-looking. No loss of substance; same appearance involves oesophagus opposite cricoid cartilage, and about four inches lower down."