and seems to help the foliage and give a better color to the bark.\*

It must be put on very thin, for if too thick it will kill the tree.

I have found the Hellebore Powder a good article to kill the worms which infest gooseberry bushes. Three years after I had placed out thirty-six plants, I picked for sale one hundred quarts, wine measure. They can be grown under fruit trees with profit. The Currant and Strawberry will also pay well. Last season I had Sharpless strawberries seven inches in circumference. These are very prolific on low land.

When referring to apples I omitted to state that I find the variety known as *King of Tompkins* does best on my soil. They are good keepers, and have been kept into July, in which month I have been shown one raised in Mahone Bay by Azariah Zwicker, Esq., in his garden, which is open to the sea winds.

Yours truly,

E. B. Hyson.

## HOW TO DESTROY INSECTS IN ORCHARDS.

From observation and past experience I find the insect enemies on the increase. Their name is legion and they call for earnest vigilance on the part of the fruit grower. In this brief article I wish to direct attention to the seven following insects which prey upon the orderd.

Canker Worm proves very destructive to the foliage. The moth appears both in spring and autumn. The females being wingless, are obliged to crawl up the stems, on which the eggs are deposited. The larva, which is ash colored or dusky brown, appears the first of June. The most effectual way of checking the ravages of this insect consists in fastening a band of coarse cloth covered with printer's ink around the trunk of the tree. The ink must be renewed every two or three days, as the dead bodies soon form a bridge across which the others could pass. Paris green and London purple dissolved in water, one part of the poison to seven hundred of water, applied with a force pump or syringe, is the most effectual remedy. If used too strong the foliage will be destroyed.

APPLE is another rings on containing they hatch them, which hatch in m inches longlateral pale of June it about ten dits eggs. along the treasily kille

The TE weave a wh branch clear torch, is the

Another better name young brance feed voracion pletely strip large as a leand can be a jar or touch position who

The Apr grey winged in great nun in the bloss eats its wa prematurely. either into t traps are set ten days, and duced in a si

<sup>\*</sup>We would advise very great caution in the use of oil. We have known trees to have been killed by the use of this unnatural remedy. An alkaline wash will prove quite as effectual and much safer.—Com.